arXiv:0805.4373v2 [math.PR] 11 Jul 2008

CONDITIONING ON AN EXTREME COMPONENT: MODEL CONSISTENCY AND REGULAR VARIATION ON CONES

BIKRAMJIT DAS AND SIDNEY I. RESNICK

Abstract. Multivariate extreme value theory assumes a multivariate domain of attraction condition for the distribution of a random vector necessitating that each component satisfy a marginal domain of attraction condition. Heffernan and Tawn (2004) and Heffernan and Resnick (2007) developed an approximation to the joint distribution of the random vector by conditioning that one of the components be extreme. The prior papers left unresolved the consistency of different models obtained by conditioning on different components being extreme and we provide understanding of this issue. We also clarify the relationship between the conditional distributions and multivariate extreme value theory. We discuss conditions under which the two models are the same and when one can extend the conditional model to the extreme value model. We also discuss the relationship between the conditional extreme value model and standard regular variation on cones of the form $[0,\infty]\times(0,\infty]$ or $(0,\infty]\times[0,\infty]$.

1. Introduction

Classical multivariate extreme value theory (abbreviated as MEVT) aims to capture the extremal dependence structure between components under a multivariate domain of attraction condition which requires that each marginal distribution belong to the domain of attraction of some univariate extreme value distribution. The theory relies on centering and scaling the components appropriately and observing the limiting behavior near the tails of the distribution. A variety of concepts have been developed in order to understand this extremal dependence structure. Multivariate extreme value theory provides a rich theory for extremal dependence in the case of asymptotic dependence (de Haan and Ferreira, 2006, de Haan and Resnick, 1977, Pickands, 1981, Resnick, 2008b) but fails to distinguish between asymptotic independence and actual independence. The extremal dependence structure in the asymptotically dependent case has been well-studied by Coles and Tawn (1991), de Haan and de Ronde (1998). The idea of *coefficient of tail dependence* developed by Ledford and Tawn (1996, 1997, 1998) provided a better understanding of asymptotically independent behavior of various components and this concept has been elaborated with the help of *hidden regular variation.* See Heffernan and Resnick (2005), Maulik and Resnick (2005), Resnick (2002, 2008a) and Resnick (2007, Chapter 8).

A different approach was provided by Heffernan and Tawn (2004) who examined multivariate distributions by conditioning on one of the components to be extreme. Their approach allowed a variety of examples of different types of asymptotic dependence and asymptotic independence. Their statistical ideas were given a more mathematical framework by Heffernan and Resnick (2007) after some slight changes in assumptions to make the theory more probabilistically viable.

Heffernan and Resnick (2007) considered a bivariate random vector (X,Y) where the distribution of Y is in the domain of attraction of an extreme value distribution G_γ, where for $\gamma\in\mathbb{R}$,

$$G_\gamma(x)=\exp\{-(1+\gamma x)^{-1/\gamma}\},\qquad 1+\gamma x>0. \tag{1.1}$$

For $\gamma=0$, the distribution function is interpreted as $G_0(x)=e^{-e^{-y}}$, $y\in\mathbb{R}$. Instead of conditioning on Y being large, their theory was developed under the equivalent assumption of the existence of a vague limit for the modified joint distribution of the a suitably scaled and centered (X,Y). Vague convergence was

---

Key words and phrases. Regular variation, domain of attraction, heavy tails, asymptotic independence, conditioned limit theory.

B. Das and S. Resnick were partially supported by ARO Contract W911NF-07-1-0078 at Cornell University.

in the space of Radon measures, $\mathbb{M}_+\big([-\infty,\infty]\times\overline{\mathbb{E}}^{(\gamma)}\big)$, and $\overline{\mathbb{E}}^{(\gamma)}, \gamma\in\mathbb{R}$, was the right closure of the set $\{x\in\mathbb{R}: 1+\gamma x>0\}$. The precise description of this vague limit is in Definition 1.1 below. It should be noted that this differs from the classical MEVT in the sense that only one of the marginal distributions is assumed to be in the domain of attraction of some univariate extreme value distribution.

In Section 2 we study the consistency issues discussed in Heffernan and Tawn (2004) for such conditional models. In practice one may have a choice of variable to condition on being large and potentially different models are therefore possible. What is the relationship between these models? We show that if conditional approximations are possible no matter which variable as the conditioning variable, then in fact the joint distribution is in a classical multivariate domain of attraction of an extreme value law. A standard case is dealt with first and later extended to a general formulation. The relationship between multivariate extreme value theory and conditioned limit theory is discussed in Section 3. Conditions under which a standardized (to be defined appropriately) regular variation model can be used in place of the conditional model are discussed in this section. We also consider conditions under which the conditional extreme value model can be extended to the multivariate extreme value model in section 4. Section 5 presents some illuminating examples to show the features of conditional models.

1.1. **Model setup and basic assumptions.**

The basic model set up for our discussion is as follows (Heffernan and Resnick, 2007):

Definition 1.1 (Conditional Extreme Value Model)**.** Suppose that for random vector $(X,Y)\in\mathbb{R}^2$, we have $Y\sim F$. We make the following assumptions:

(1) F is in the domain of attraction of an extreme value distribution, G_γ for some $\gamma\in\mathbb{R}$, as defined in (1.1); that is, there exist functions $a(t)>0, b(t)\in\mathbb{R}$ such that, as $t\to\infty$,

$$t(1-F(a(t)y+b(t)))=t\mathbf{P}\left(\frac{Y-b(t)}{a(t)}>y\right)\to(1+\gamma y)^{-1/\gamma},\qquad 1+\gamma y>0. \tag{1.2}$$

(2) There exist functions $\alpha(t)>0$ and $\beta(t)\in\mathbb{R}$ and a non-null Radon measure μ on Borel subsets of $[-\infty,\infty]\times\overline{\mathbb{E}}^{(\gamma)}$ such that for each $y\in\mathbb{E}^{(\gamma)}$,

$$[a]\quad t\mathbf{P}\left(\frac{X-\beta(t)}{\alpha(t)}\le x, \frac{Y-b(t)}{a(t)}>y\right)\to\mu([-\infty,x]\times(y,\infty]),\qquad \text{as } t\to\infty, \tag{1.3}$$

for (x,y) continuity points of the limit.

$$[b]\quad \mu([-\infty,x]\times(y,\infty])\quad \text{is not a degenerate distribution in } x, \tag{1.4}$$

$$[c]\quad \mu([-\infty,x]\times(y,\infty])<\infty. \tag{1.5}$$

$$[d]\quad H(x):=\mu([-\infty,x]\times(0,\infty]) \text{ is a probability distribution.} \tag{1.6}$$

A non-null Radon measure $\mu(\cdot)$ is said to satisfy the *conditional non-degeneracy conditions* if both of (1.4) and (1.5) hold. We will say that (X,Y) follows a *conditional extreme value model* (abbreviated as CEVM) if conditions (1) and (2) above are satisfied. The reason for this name is that (1.3),(1.4) and (1.5) imply that for continuity points (x,y) of $\mu(\cdot)$,

$$\mathbf{P}\left(\frac{X-\beta(t)}{\alpha(t)}\le x\,\middle|\, Y>b(t)\right)\to H(x)=\mu([-\infty,x]\times(0,\infty]).\qquad (t\to\infty) \tag{1.7}$$

Also note that (1.3) can be viewed in terms of vague convergence of measures in $\mathbb{M}_+\big([-\infty,\infty]\times\overline{\mathbb{E}}^{(\gamma)}\big)$.

Under the above assumptions Heffernan and Resnick (2007) use a convergence to types argument to get properties of the normalizing and centering functions: there exists functions $\psi_1(\cdot),\psi_2(\cdot)$ such that

$$\lim_{t\to\infty}\frac{\alpha(tc)}{\alpha(t)}=\psi_1(c),\qquad\qquad \lim_{t\to\infty}\frac{\beta(tc)-\beta(t)}{\alpha(t)}=\psi_2(c). \tag{1.8}$$

This implies that $\psi(c) = c^{\rho}$ for some $\rho \in \mathbb{R}$ (de Haan and Ferreira, 2006, Theorem B.1.3). ψ_2 can be either 0 or $\psi_2(c) = k\frac{c^{\rho}-1}{\rho}$ for some $c \neq 0$ (de Haan and Ferreira, 2006, Theorem B.2.1). We refer often to these properties.

How different is the CEVM from classical MEVT model? Under what circumstances can we extend from assumptions 1 and 2 to multivariate extreme value theory? We deal with questions like these in the following sections.

1.2. **Notation.** We list below commonly used notation. The Appendix in Section 6 contains information on regularly varying functions and extensions to such things as Π-varying functions, as well as a rapid review of vague convergence.

$\mathbb{R}_+^d$	$[0,\infty)^d$.
$\overline{\mathbb{R}}_+^d$	$[0,\infty]^d$. Also denote similarly $\overline{\mathbb{R}}^d = [-\infty,\infty]^d$.
$\mathbb{E}^*$	A nice subset of the compactified finite dimensional Euclidean space. Often denoted $\mathbb{E}$ with different subscripts and superscripts as required.
$\mathcal{E}^*$	The Borel $\sigma-$field of the subspace $\mathbb{E}^*$.
$\mathbb{M}_+(\mathbb{E}^*)$	The class of Radon measures on Borel subsets of $\mathbb{E}^*$.
$f^{\leftarrow}$	The left continuous inverse of a monotone function f. For an increasing function $f^{\leftarrow}(x) = \inf\{y : f(y) \geq x\}$. For a decreasing function $f^{\leftarrow}(x) = \inf\{y : f(y) \leq x\}$.
RV_{ρ}	The class of regularly varying functions with index ρ defined in (6.1).
Π	The function class Π reviewed in Section 6.1 along with subclasses $\Pi_+\big(a(\cdot)\big)$ and $\Pi_-\big(a(\cdot)\big)$ and *auxiliary function* $a(\cdot)$.
$\xrightarrow{v}$	Vague convergence of measures; see Section 6.3.
G_{γ}	An extreme value distribution given by (1.1), with parameter $\gamma \in \mathbb{R}$, in the Von Mises parameterization.
$\mathbb{E}^{(\gamma)}$	$\{x : 1+\gamma x > 0\}$ for $\gamma \in \mathbb{R}$.
$\overline{\mathbb{E}}^{(\gamma)}$	The closure on the right of the interval $\mathbb{E}^{(\gamma)}$.
$\overline{\overline{\mathbb{E}}}^{(\gamma)}$	The closure on both sides of the interval $\mathbb{E}^{(\gamma)}$.
$\mathbb{E}^{(\lambda,\gamma)}$	$\overline{\overline{\mathbb{E}}}^{(\lambda)} \times \overline{\overline{\mathbb{E}}}^{(\gamma)} \setminus \{(-\frac{1}{\lambda}, -\frac{1}{\gamma})\}$.
$D(G_{\gamma})$	The domain of attraction of the extreme value distribution G_{γ}; i.e., the set of F's satisfying (1.2). For $\gamma > 0$, $F \in D(G_{\gamma})$ is equivalent to $1 - F \in RV_{1/\gamma}$.

2. Consistency of Conditional Extreme Value Models

Suppose that the $(X,Y) \in \mathbb{R}^2$ satisfying conditions (1.2) to (1.6) also satisfies the following conditions. We have $X \sim H$ where H is in the domain of attraction of an extreme value distribution and we have a convergence assumption similar to assumptions (1.3) to (1.6) for the joint distribution of (X,Y) where X is large. One ponders whether this would imply that (X,Y) is jointly in the domain of attraction of a multivariate extreme value distribution. The answer is yes.

2.1. **Consistency: the standard case.** Let us start with a simple case. Assume for (X,Y) as mentioned in the previous paragraph, the centering functions are all *zero* and the norming functions are *identity functions.*

Now set

$$\begin{aligned} \mathbb{E} &= [0,\infty]^2 \setminus \{\mathbf{0}\}, & \mathbb{E}_0 &= (0,\infty] \times (0,\infty], \\ \mathbb{E}_{\sqcap} &= [0,\infty] \times (0,\infty], & \mathbb{E}_{\sqsupset} &= (0,\infty] \times [0,\infty]. \end{aligned}$$

Figure 1 illustrates these four types of cones in two dimensions. Before proceeding, we review the definition of multivariate regular variation on cones (Resnick (2007, page 173), Davydov et al. (2007), Resnick (2008a).



FIGURE 1. The different cones in 2-dimensions

Definition 2.1. $\mathfrak{C} \subset \overline{\mathbb{R}}^d$ is said to be a cone if $x \in \mathfrak{C}$ implies that $tx \in \mathfrak{C}$ for any $t > 0$. Now a $d-$dimensional random vector $\boldsymbol{Z} \in \mathbb{R}^d$ is said to be *multivariate regularly varying* on cone $\mathfrak{C}$ in $\overline{\mathbb{R}}^d$ if there exists a function $b(t) \to \infty$ and a non-null Radon measure ν on $\mathfrak{C}$ such that, as $t \to \infty$,

$$t\mathbf{P}\left[\frac{\boldsymbol{Z}}{b(t)} \in \cdot\right] \xrightarrow{v} \nu(\cdot), \qquad \text{in } \mathbb{M}_+(\mathfrak{C}). \tag{2.1}$$

Remark 2.1. It follows from (2.1) that the limit measure $\nu(\cdot)$ satisfies the homogeneity property that for a relatively compact set $B \subset \mathfrak{C}$,

$$\nu(cB) = c^{-\alpha}\nu(B) \qquad c > 0, \tag{2.2}$$

for some $\alpha > 0$. This also implies that $b(\cdot)$ is regularly varying with index $1/\alpha$.

Remark 2.2. The regular variation in Definition 2.1 is said to be *standard* if $\boldsymbol{Z} \in \mathbb{R}_+^d$ and $b(t) \equiv t$. In this case we have equation (2.2) with $\alpha = 1$.

Note that $\mathbb{E}, \mathbb{E}_{\sqcap}, \mathbb{E}_{\sqsupset}, \mathbb{E}_0$ are all cones in $\overline{\mathbb{R}}^2$. The following result shows that standard regular variation on both $\mathbb{E}_{\sqcap}$ and $\mathbb{E}_{\sqsupset}$ imply standard regular variation on the bigger cone $\mathbb{E}_{\sqcap} \cup \mathbb{E}_{\sqsupset} = \mathbb{E}$. This is the introduction to the more general consistency results in the CEVM.

Theorem 2.1. *Suppose we have a bivariate random vector* $(X, Y) \in \mathbb{R}_+^2$*. Now assume that*

$$t\mathbf{P}\Big[\Big(\frac{X}{t}, \frac{Y}{t}\Big) \in \cdot\Big] \xrightarrow{v} \mu(\cdot) \ \text{in } \mathbb{M}_+\big(\mathbb{E}_\sqcap\big), \tag{2.3}$$

$$t\mathbf{P}\Big[\Big(\frac{X}{t}, \frac{Y}{t}\Big) \in \cdot\Big] \xrightarrow{v} \nu(\cdot) \ \text{in } \mathbb{M}_+\big(\mathbb{E}_\sqsupset\big). \tag{2.4}$$

where both μ *and* ν *satisfy appropriate conditional non-degeneracy conditions corresponding to* (1.4)–(1.6)*. Then* (X, Y) *is standard regularly varying in* $\mathbb{E}$*, i.e.,*

$$t\mathbf{P}\Big[\Big(\frac{X}{t}, \frac{Y}{t}\Big) \in \cdot\Big] \xrightarrow{v} (\mu \diamond \nu)(\cdot) \ \text{in } \mathbb{M}_+\big(\mathbb{E}\big) \tag{2.5}$$

where $(\mu \diamond \nu)$ *is a Radon measure on* $\mathbb{E}$ *such that*

$$(\mu \diamond \nu)|_{\mathbb{E}_\sqcap}(\cdot) = \mu(\cdot) \ \text{ on } \mathbb{E}_\sqcap \qquad \text{and} \qquad (\mu \diamond \nu)|_{\mathbb{E}_\sqsupset}(\cdot) = \nu(\cdot) \ \text{ on } \mathbb{E}_\sqsupset.$$

The proof of this theorem is provided in Section 6.4.1 since a more general result is stated and proved next. Note, however, that the proof for Theorem 2.1 is relatively easy because of the standard case assumptions and is instructive to read.

2.2. **Consistency: the general case.** Thus we see that multivariate regular variation on both the cones $\mathbb{E}_\sqsupset$ and $\mathbb{E}_\sqcap$ implies multivariate regular variation on the larger cone $\mathbb{E}_\sqsupset \cup \mathbb{E}_\sqcap = \mathbb{E}$. Now we will discuss the general situation in which each marginal distribution is in the domain of attraction of an extreme value distribution. Recall our notation, $\mathbb{E}^{(\gamma)} = \{x \in \mathbb{R} : 1 + \gamma x > 0\}$ for $\gamma \in \mathbb{R}$. We denote $\overline{\mathbb{E}}^{(\gamma)}$ to be the right closure of $\mathbb{E}^{(\gamma)}$, i.e.,

$$\overline{\mathbb{E}}^{(\gamma)} = \begin{cases} (-\frac{1}{\gamma}, \infty] & \gamma > 0 \\ (-\infty, \infty] & \gamma = 0 \\ (-\infty, -\frac{1}{\gamma}] & \gamma < 0. \end{cases} \tag{2.6}$$

$\overline{\overline{\mathbb{E}}}^{(\gamma)}$ denotes the set we get by closing $\mathbb{E}^{(\gamma)}$ on both sides. Also denote $\mathbb{E}^{(\lambda,\gamma)} := \overline{\overline{\mathbb{E}}}^{(\lambda)} \times \overline{\overline{\mathbb{E}}}^{(\gamma)} \setminus \{(-\frac{1}{\lambda}, -\frac{1}{\gamma})\}$.

Theorem 2.2. *Suppose we have a bivariate random vector* $(X, Y) \in \mathbb{R}^2$ *and non-negative functions* $\alpha(\cdot), a(\cdot)$*,* $\chi(\cdot), c(\cdot)$ *and real valued functions* $\beta(\cdot), b(\cdot), \phi(\cdot), d(\cdot)$ *such that*

$$t\mathbf{P}\Big[\Big(\frac{X - \beta(t)}{\alpha(t)}, \frac{Y - b(t)}{a(t)}\Big) \in \cdot\Big] \xrightarrow{v} \mu(\cdot) \ \text{in } \mathbb{M}_+\big([-\infty, \infty] \times \overline{\mathbb{E}}^{(\gamma)}\big), \tag{2.7}$$

$$t\mathbf{P}\Big[\Big(\frac{X - \phi(t)}{\chi(t)}, \frac{Y - d(t)}{c(t)}\Big) \in \cdot\Big] \xrightarrow{v} \nu(\cdot) \ \text{in } \mathbb{M}_+\big(\overline{\mathbb{E}}^{(\lambda)} \times [-\infty, \infty]\big) \tag{2.8}$$

for some $\lambda, \gamma \in \mathbb{R}$*, where both* μ *and* ν *satisfy the appropriate conditional non-degeneracy conditions corresponding to* (1.4) *and* (1.5)*. Then* (X, Y) *is in the domain of attraction of a multivariate extreme value distribution on* $\mathbb{E}^{(\lambda,\gamma)}$ *in the following sense:*

$$t\mathbf{P}\Big[\Big(\frac{X - \phi(t)}{\chi(t)}, \frac{Y - b(t)}{a(t)}\Big) \in \cdot\Big] \xrightarrow{v} (\mu \diamond \nu)(\cdot) \ \text{in } \mathbb{M}_+\big(\mathbb{E}^{(\lambda,\gamma)}\big)$$

where $(\mu \diamond \nu)(\cdot)$ *is a non-null Radon measure on* $\mathbb{E}^{(\lambda,\gamma)}$*.*

Proof. Let us assume that $\lambda > 0, \gamma > 0$ first. The other cases can be dealt with similarly. It should be noted now, that (2.7) and (2.8) respectively imply that

$$t\mathbf{P}\Big(\frac{Y - b(t)}{a(t)} > y\Big) \to (1 + \gamma y)^{-1/\gamma}, \ 1 + \gamma y > 0, \tag{2.9}$$

$$t\mathbf{P}\Big(\frac{X - \phi(t)}{\chi(t)} > x\Big) \to (1 + \lambda x)^{-1/\lambda}, \ 1 + \lambda x > 0. \tag{2.10}$$

Hence for $(x,y) \in \mathbb{E}^{(\lambda)} \times \mathbb{E}^{(\gamma)}$,which are continuity points of the limit measures μ and ν,

$$\begin{aligned} Q_t(x,y) :&= t\mathbf{P}\Big[\Big(\frac{X-\phi(t)}{\chi(t)}, \frac{Y-b(t)}{a(t)}\Big) \in \big([-\infty,x]\times[-\infty,y]\big)^c\Big] \\ &= t\mathbf{P}\Big[\frac{X-\phi(t)}{\chi(t)} > x\Big] + t\mathbf{P}\Big[\frac{Y-b(t)}{a(t)} > y\Big] - t\mathbf{P}\Big[\frac{X-\phi(t)}{\chi(t)} > x, \frac{Y-b(t)}{a(t)} > y\Big] \\ &= A_t(x) + B_t(y) + C_t(x,y) \qquad \text{(say).} \end{aligned} \tag{2.11}$$

If we can show that $Q_t(x,y)$ has a limit and the limit is non-degenerate over (x,y) then we are done. As $t \to \infty$ we have the limits for $A_t(x)$ and $B_t(y)$ from equations (2.10) and (2.9) respectively. Clearly $0 \le C_t(x,y) \le \min(A_t(x), B_t(y))$ and these inequalities would hold for any limit of Q_t as well.

From (Heffernan and Resnick, 2007, Proposition 1), there exist functions $\psi_1(\cdot), \psi_2(\cdot), \psi_3(\cdot), \psi_4(\cdot)$ such that for $z > 0$,

$$\lim_{t\to\infty}\frac{\alpha(tz)}{\alpha(t)} = \psi_1(z) = z^{\rho_1}, \qquad \lim_{t\to\infty}\frac{\beta(tz)-\beta(t)}{\alpha(t)} = \psi_2(z), \tag{2.12}$$

$$\lim_{t\to\infty}\frac{c(tz)}{c(t)} = \psi_3(z) = z^{\rho_2}, \qquad \lim_{t\to\infty}\frac{d(tz)-d(t)}{c(t)} = \psi_4(z). \tag{2.13}$$

for some real constants ρ_1 and ρ_2. Assume ρ_1 and ρ_2 to be positive for the time being. Here either $\psi_2(z) = 0$ which would imply $\lim\limits_{t\to\infty}\frac{\beta(t)}{\alpha(t)} = 0$ (from Bingham et al. (1987, Theorem 3.1.12 a,c)) or we can have $\psi_2(z) = k\frac{z^{\rho_1}-1}{\rho_1}$ for some $k \neq 0$, which means $\lim\limits_{t\to\infty}\frac{\beta(t)}{\alpha(t)} = \frac{k}{\rho_1}$ (de Haan and Ferreira (2006, Proposition B.2.2)). Hence allowing the constant k to be zero as well, we can write both cases as $\lim\limits_{t\to\infty}\frac{\beta(t)}{\alpha(t)} = \frac{k_1}{\rho_1}$ for some $k_1 \in \mathbb{R}$. Similarly we have $\lim\limits_{t\to\infty}\frac{d(t)}{c(t)} = \frac{k_2}{\rho_2}$ for some $k_2 \in \mathbb{R}$.

Additionally, from the marginal domain of attraction conditions for X, Y we have

$$\lim_{t\to\infty}\frac{b(tz)-b(t)}{a(t)} = \frac{z^\gamma-1}{\gamma}, \quad \text{for } y>0, \quad \text{which implies } \lim_{t\to\infty}\frac{a(tz)}{a(t)} = z^\gamma, \tag{2.14}$$

$$\text{and } \lim_{t\to\infty}\frac{\phi(tw)-\phi(t)}{\chi(t)} = \frac{w^\lambda-1}{\lambda}, \quad \text{for } w>0, \quad \text{which implies } \lim_{t\to\infty}\frac{\chi(tw)}{\chi(t)} = w^\lambda. \tag{2.15}$$

Observe that

$$\begin{aligned} C_t(x,y) &= t\mathbf{P}\Big[\frac{X-\phi(t)}{\chi(t)} > x, \frac{Y-b(t)}{a(t)} > y\Big] \\ &= t\mathbf{P}\Big[\frac{X-\beta(t)}{\alpha(t)} > \Big(x+\frac{\phi(t)}{\chi(t)}\Big)\frac{\chi(t)}{\alpha(t)} - \frac{\beta(t)}{\alpha(t)}, \frac{Y-b(t)}{a(t)} > y\Big]. \end{aligned} \tag{2.16}$$

We can also write

$$C_t(x,y) = t\mathbf{P}\Big[\frac{X-\phi(t)}{\chi(t)} > x, \frac{Y-d(t)}{c(t)} > \Big(y+\frac{b(t)}{a(t)}\Big)\frac{a(t)}{c(t)} - \frac{d(t)}{c(t)}\Big] \tag{2.17}$$

From de Haan and Ferreira (2006, Proposition B.2.2) we have that

$$\frac{b(t)}{a(t)} \to \frac{1}{\gamma} \quad \text{and} \quad \frac{\phi(t)}{\chi(t)} \to \frac{1}{\lambda}. \tag{2.18}$$

We analyze $C_t(x,y)$ for the different cases now. First we will show that at least one of the limits $\lim\limits_{t\to\infty}\frac{\chi(t)}{\alpha(t)}$ and $\lim\limits_{t\to\infty}\frac{a(t)}{c(t)}$ has to exist. Suppose both do not exist. We have for $(x,y) \in \mathbb{E}^{(\lambda)} \times \mathbb{E}^{(\gamma)}$, which are continuity points of the limit measures μ and ν,

$$t\mathbf{P}\Big[\frac{X-\beta(t)}{\alpha(t)} > x, \frac{Y-b(t)}{a(t)} > y\Big] \to \mu((x,\infty]\times(y,\infty]), \tag{2.19}$$

$$t\mathbf{P}\Big[\frac{X-\phi(t)}{\chi(t)}>x,\frac{Y-d(t)}{c(t)}>y\Big]\to\nu((x,\infty]\times(y,\infty]). \tag{2.20}$$

Now (2.19) implies that

$$t\mathbf{P}\Big[\frac{X-\phi(t)}{\chi(t)}\frac{\chi(t)}{\alpha(t)}+\frac{\phi(t)-\beta(t)}{\alpha(t)}>x,\frac{Y-d(t)}{c(t)}\frac{c(t)}{a(t)}+\frac{d(t)-b(t)}{a(t)}>y\Big]\to\mu\big((x,\infty]\times(y,\infty]\big)$$

which is equivalent to

$$t\mathbf{P}\Big[\frac{X-\phi(t)}{\chi(t)}>\frac{\alpha(t)}{\chi(t)}\big(x-\frac{\phi(t)-\beta(t)}{\alpha(t)}\big),\frac{Y-d(t)}{c(t)}>\frac{a(t)}{c(t)}\big(y-\frac{d(t)-b(t)}{a(t)}\big)\Big]\to\mu\big((x,\infty]\times(y,\infty]\big).$$

From (2.20) we also have that the left side of the previous line has a limit

$$t\mathbf{P}\Big[\frac{X-\phi(t)}{\chi(t)}>\frac{\alpha(t)}{\chi(t)}\big(x-\frac{\phi(t)-\beta(t)}{\alpha(t)}\big),\frac{Y-d(t)}{c(t)}>\frac{a(t)}{c(t)}\big(y-\frac{d(t)-b(t)}{a(t)}\big)\Big]$$
$$\to\nu\big((f(x),\infty]\times(g(y),\infty]\big)$$

for some $(f(x),g(y))$, assumed to be a continuity point of the limit ν, iff as $t\to\infty$, the following two limits hold,

$$\frac{\alpha(t)}{\chi(t)}\Big(x-\frac{\phi(t)-\beta(t)}{\alpha(t)}\Big)\to f(x), \tag{2.21}$$

$$\frac{a(t)}{c(t)}\Big(y-\frac{d(t)-b(t)}{a(t)}\Big)\to g(y). \tag{2.22}$$

For ν to be non-degenerate f and g should be non-constant and we also have $\mu\big((x,\infty]\times(y,\infty]\big)=\nu\big((f(x),\infty]\times(g(y),\infty]\big)$. Considering (2.21) and (2.22) we can see that the limit as $t\to\infty$ exists if and only if $\lim_{t\to\infty}\frac{a(t)}{c(t)}$ and $\lim_{t\to\infty}\frac{\chi(t)}{\alpha(t)}$ exists.

We conclude $\lim_{t\to\infty}\frac{\chi(t)}{\alpha(t)}\in[0,\infty]$ and consider the following cases.

- **Case 1:** $\lim_{t\to\infty}\frac{\chi(t)}{\alpha(t)}=\infty$.
  Consider (2.16) and note
  $$\Big(x+\frac{\phi(t)}{\chi(t)}\Big)\frac{\chi(t)}{\alpha(t)}-\frac{\beta(t)}{\alpha(t)}\to\Big(x+\frac{1}{\lambda}\Big)\times\infty-\frac{k_1}{\rho_1}=\infty,$$
  which entails
  $$\lim_{t\to\infty}C_t(x,y)=\mu(\{\infty\}\times(y,\infty])=0.$$
  Hence
  $$\lim_{t\to\infty}Q_t(x,y)=(1+\lambda x)^{-1/\lambda}+(1+\gamma y)^{-1/\gamma}.$$
- **Case 2:** $\lim_{t\to\infty}\frac{\chi(t)}{\alpha(t)}=M\in(0,\infty)$.
  Again from (2.16), we have
  $$\Big(x+\frac{\phi(t)}{\chi(t)}\Big)\frac{\chi(t)}{\alpha(t)}-\frac{\beta(t)}{\alpha(t)}\to\Big(x+\frac{1}{\lambda}\Big)\times M-\frac{k_1}{\rho_1}=f(x),\quad\text{(say)}.$$
  Therefore
  $$\lim_{t\to\infty}C_t(x,y)=\mu((f(x),\infty]\times(y,\infty])\le(1+\lambda y)^{-1/\lambda}$$
  with strict inequality holding for some x because of the non-degeneracy condition (1.4) for μ. Hence
  $$\lim_{t\to\infty}Q_t(x,y)=((1+\lambda x)^{-1/\lambda}+(1+\gamma y)^{-1/\gamma}-\mu((f(x),\infty]\times(y,\infty]).$$

- **Case 3:** $\lim_{t\to\infty} \frac{\chi(t)}{\alpha(t)} = 0$.

  In this case (2.16) leads to a degenerate limit in x for $C_t(x,y)$ and putting $M_1 = \frac{k}{\rho_1}$ we get

  $$\lim_{t\to\infty} C_t(x,y) = \mu((M_1,\infty] \times (y,\infty]) =: f_1(y) \le (1+\gamma y)^{-1/\gamma}.$$

  So consider (2.17).

  (1) If $\lim_{t\to\infty} \frac{a(t)}{c(t)}$ exists in $(0,\infty]$, then we can use a similar technique as in case 1 or case 2 to obtain a non-degenerate limit for $Q_t(x,y)$.

  (2) Assume $\lim_{t\to\infty} \frac{a(t)}{c(t)} = 0$. Then for some $M_2 \in \mathbb{R}$,

  $$\lim_{t\to\infty} C_t(x,y) = \nu((x,\infty] \times (M_2,\infty] =: f_2(x) \le (1+\lambda x)^{-1/\lambda}$$

  Therefore we have for any $(x,y) \in \mathbb{E}^{(\lambda)} \times \mathbb{E}^{(\gamma)}$,which are continuity points of the limit measures μ and ν,

  $$f_1(y) = \mu((M_1,\infty] \times (y,\infty]) = \nu((x,\infty] \times (M_2,\infty]) = f_2(x).$$

  It is easy to check now that for any $(x,y) \in \mathbb{E}^{(\lambda)} \times \mathbb{E}^{(\gamma)}$, which are continuity points of the limit measures μ and ν, we have $f_1(y) = f_2(x) = 0$. Hence $C_t(x,y) \to 0$ and thus $Q_t(x,y)$ has a non-degenerate limit.

This proves the result. □

2.3. **Consistency: the absolutely continuous case.** It is instructive to consider the consistency issue when (X,Y) has a joint density since calculations become more explicit. We state the consistency result particularizing Theorem 2.2; the proof for the standard case is provided in Section 6.4.2.

Proposition 2.3. *Assume g_ρ denotes the density of an extreme value distribution G_ρ with shape parameter $\rho \in \mathbb{R}$ in* (1.1) *and that $(X,Y) \in \mathbb{R}^2$ is a bivariate random vector. We suppose*

1. *(X,Y) has a density $f_{X,Y}(x,y)$.*
2. *The marginal densities f_X, f_Y satisfy:*

$$t\alpha(t) f_X(\alpha(t)x+\beta(t)) \to g_\lambda(x), x \in \mathbb{E}^{(\lambda)}, \qquad t a(t) f_Y(a(t)y+b(t)) \to g_\gamma(y), y \in \mathbb{E}^{(\gamma)}. \tag{2.23}$$

3. *The joint density satisfies*

$$t\alpha(t)a(t) f_{X,Y}(\alpha(t)x+\beta(t), a(t)y+b(t)) \to g_1(x,y) \in L^1([-\infty,\infty] \times \overline{\mathbb{E}}^{(\gamma)}), \tag{2.24}$$

$$t\alpha(t)a(t) f_{X,Y}(\alpha(t)x+\beta(t), a(t)y+b(t)) \to g_2(x,y) \in L^1(\overline{\mathbb{E}}^{(\lambda)} \times [-\infty,\infty]), \tag{2.25}$$

   where $g_1(x,y), g_2(x,y) \ge 0$ are non-trivial, 0 outside of $[-\infty,\infty] \times \overline{\mathbb{E}}^{(\gamma)}$ and $\overline{\mathbb{E}}^{(\lambda)} \times [-\infty,\infty]$ respectively, and we have

   (a) *$g_\gamma^{-1}(v) g_1(u,v)$ is a probability density in v for each $u > 0$,*

   (b) *$g_\lambda^{-1}(u) g_2(u,v)$ is a probability density in u for each $v > 0$.*

Then

$$t\mathbf{P}\Big[\Big(\frac{X-\beta(t)}{\alpha(t)}, \frac{Y-b(t)}{a(t)}\Big) \in \cdot\Big] \xrightarrow{v} \mu(\cdot) \text{ in } \mathbb{M}_+\big(\mathbb{E}^{(\lambda,\gamma)}\big),$$

where

$$\mu(A) = \int\limits_{\{(u,v)\in A\}} g(u,v)\,du\,dv \qquad \text{for } A \in \mathcal{B}\big(\mathbb{E}^{(\lambda,\gamma)}\big),$$

and $g(x,y) = g_1(x,y) \vee g_2(x,y)$.

3. Connecting Standard Regular Variation to the Conditional Extreme Value Model

We have seen in the previous sections that questions about the general conditional model are effectively analyzed by starting with standard regular variation on our special cones ($\mathbb{E}_{\sqsupset}$ or $\mathbb{E}_{\sqcap}$). A pertinent question to ask here is, whether standardization of the conditional extreme value model is always possible. A partial answer has been provided in Heffernan and Resnick (2007, Section 2.4). We consider this issue in more detail in this section. We start by making precise what we mean by *standardization*.

3.1. **Standardization.**

Standardization is the process of marginally transforming a random vector $\boldsymbol{X}$ into a different vector $\boldsymbol{Z}^*$, $\boldsymbol{X} \mapsto \boldsymbol{Z}^*$, so that the distribution of $\boldsymbol{Z}^*$ is standard regularly varying on a cone $\mathbb{E}^*$. For some Radon measure $\mu^*(\cdot)$

$$tP\Big[\frac{\boldsymbol{Z}^*}{t} \in \cdot\,\Big] \xrightarrow{v} \mu^*(\cdot), \qquad \text{in } \mathbb{M}_+(\mathbb{E}^*).$$

In general, depending on the cone, this says one or more components of $\boldsymbol{Z}^*$ are asymptotically Pareto. For the classical multivariate extreme value theory case, each is asymptotically Pareto and then $\mathbb{E}^* = \mathbb{E} = [\boldsymbol{0}, \boldsymbol{\infty}] \setminus \{\boldsymbol{0}\}$. The technique is used in classical multivariate extreme value theory to characterize multivariate domains of attraction and dates at least to de Haan and Resnick (1977). See also Resnick (2008b, Chapter 5), de Haan and Ferreira (2006), Mikosch (2005, 2006), Resnick (2007).

3.1.1. *Theoretical advantages of standardization:*

- Standardization is analogous to the copula transformation but is better suited to studying limit relations (Klüppelberg and Resnick, 2008).
- In Cartesian coordinates, the limit measure has scaling property:
$$\mu^*(c\,\cdot\,) = c^{-1}\mu^*(\cdot), \quad c > 0.$$
- The scaling in Cartesian coordinates allows transformation to polar coordinates to yield a product measure: An angular measure exists allowing characterization of limits:
$$\mu^*\{\boldsymbol{x} : \|\boldsymbol{x}\| > r, \frac{\boldsymbol{x}}{\|\boldsymbol{x}\|} \in \Lambda\} = r^{-1}S(\Lambda),$$
for Borel subsets Λ of the unit sphere in $\mathbb{E}^*$.

Note that for classical multivariate extreme value theory, S is a finite measure which we may take to be a probability measure without loss of generality. However, when $\mathbb{E}^* = \mathbb{E}_{\sqcap}$, S is NOT necessarily finite. This is because absence of the horizontal axis boundary in $\mathbb{E}_{\sqcap}$ implies the unit sphere is not compact.

3.1.2. *Standardizing functions.* The most useful circumstance for standardization is discussed in the following definition.

Definition 3.1. Suppose $\boldsymbol{X} = (X_1, X_2, \dots, X_d)$ is a vector-valued random variable in $\mathbb{R}^d$ which satisfies:

$$t\mathbf{P}\Big[\Big(\frac{X_1 - \beta_1(t)}{\alpha_1(t)}, \frac{X_2 - \beta_2(t)}{\alpha_2(t)}, \dots, \frac{X_d - \beta_d(t)}{\alpha_d(t)}\Big) \in \cdot\,\Big] \xrightarrow{v} \mu(\cdot) \qquad \text{on } \mathbb{M}_+(\mathfrak{D}), \tag{3.1}$$

for some $\mathfrak{D} \subset \overline{\mathbb{R}}^d$, $\alpha_i(t) > 0, \beta_i(t) \in \mathbb{R}$, for $i = 1, \dots, d$. Suppose we have $\boldsymbol{f} = (f_1, \dots, f_d)$ such that, for $i = 1, \dots, d$:

(a) f_i : Range of $X_i \to (0, \infty)$,
(b) f_i is monotone,
(c) $\nexists K > 0$ such that $|f_i| \le K$.

Then $\boldsymbol{f}$ *standardizes* $\boldsymbol{X}$ if $\boldsymbol{Z}^* = \boldsymbol{f}(\boldsymbol{X})$ satisfies

$$t\mathbf{P}\Big[\Big(\frac{\boldsymbol{Z}^*}{t} \in \cdot\,\Big] = t\mathbf{P}\Big[\Big(\frac{f_1(X_1)}{t}, \frac{f_2(X_2)}{t}, \dots, \frac{f_d(X_d)}{t}\Big) \in \cdot\,\Big] \xrightarrow{v} \mu^*(\cdot) \text{ on } \mathbb{M}_+(\mathbb{E}^*), \tag{3.2}$$

where $\mathbb{E}^*$ is some cone in $\overline{\mathbb{R}}_+^d$, and μ^* is Radon. Call $\boldsymbol{f}$ the *standardizing function* and say (3.2) is the *standardization* of (3.1).

For the conditional model defined in Definition 1.1 in Section 1.1 where F, the distribution of Y, satisfies $F \in D(G_1)$, we can always standardize Y with

$$b^{\leftarrow}(\cdot) = \Big(\frac{1}{1-F}\Big)^{\leftarrow}(\cdot).$$

We define $Y^* = b^{\leftarrow}(Y)$ to be the standardized version of Y and the standardizing function is $b^{\leftarrow}$. See Heffernan and Resnick (2007).

3.2. **When can the conditional extreme value model be standardized?** Suppose Definition 1.1 holds; that is

$$t\mathbf{P}\Big[\Big(\frac{X-\beta(t)}{\alpha(t)}, \frac{Y-b(t)}{a(t)}\Big) \in \cdot\Big] \xrightarrow{v} \mu(\cdot) \qquad \text{in } \mathbb{M}_+\big([-\infty,\infty] \times \overline{\mathbb{E}}^{(\gamma)}\big).$$

Heffernan and Resnick (2007) show that standardization in the above equation is possible unless $(\psi_1, \psi_2) = (1,0)$ which is equivalent to the limit measure being a product measure. We show that the converse is true too. So when the limit measure is not a product measure, we can always reduce to standard regular variation on a cone $\mathbb{E}_{\sqcap}$, and conversely we can think of the general conditional model as a transformation of standard regular variation on $\mathbb{E}_{\sqcap}$.

We begin with initial results about the impossibility of the limit measure being a product in the standardized convergence on $\mathbb{E}_{\sqcap}$, gradually leading to our final result in Proposition 3.3.

Lemma 3.1. *Suppose (X,Y) is standard regularly varying on the cone $\mathbb{E}_{\sqcap}$, such that,*

$$t\mathbf{P}\Big[\Big(\frac{X}{t}, \frac{Y}{t}\Big) \in \cdot\Big] \xrightarrow{v} \mu(\cdot) \qquad \text{in } \mathbb{M}_+\big(\mathbb{E}_{\sqcap}\big) \tag{3.3}$$

for some non-null Radon measure $\mu(\cdot)$ on $\mathbb{E}_{\sqcap}$, satisfying the conditional non-degeneracy conditions as in (1.4) *and* (1.5)*. Then $\mu(\cdot)$ cannot be a product measure.*

Proof. If μ is a product measure we have

$$\mu\big([0,x] \times (y,\infty]\big) = G(x)y^{-1} \quad \text{for } x \geq 0, y > 0 \tag{3.4}$$

for some finite distribution function G on $[0,\infty)$. Now (3.3) implies that μ is homogeneous of order -1, i.e.,

$$\mu(c\Lambda) = c^{-1}\mu(\Lambda), \qquad \forall c > 0, \tag{3.5}$$

where Λ is a Borel subset of $\mathbb{E}_{\sqcap}$. Therefore

$$\begin{aligned} \mu\big(c([0,x] \times (y,\infty])\big) &= \mu\big([0,cx] \times (cy,\infty]\big) \\ &= G(cx) \times \frac{1}{cy}, \qquad \text{using (3.4)} \\ &= c^{-1}G(cx)y^{-1}. \end{aligned}$$

Moreover,

$$\mu\big(c([0,x] \times (y,\infty])\big) = c^{-1}G(x)y^{-1}, \qquad \text{using (3.4) and (3.5).}$$

Therefore

$$G(cx) = G(x) \quad \forall c > 0, x > 0.$$

Hence for fixed $y \in \mathbb{E}^{(\gamma)}, c > 0, x > 0$,

$$\mu\big([0,cx] \times (y,\infty]\big) = G(cx)y^{-1} = G(x)y^{-1} = \mu\big([0,x] \times (y,\infty]\big).$$

Therefore μ becomes a degenerate distribution in x, contradicting our conditional non-degeneracy assumptions. Thus $\mu(\cdot)$ cannot be a product measure. ☐

Lemma 3.1 means that standard regular variation on $\mathbb{E}_\sqcap$ with a limit measure satisfying the conditional non-degeneracy conditions implies that the limit cannot be a product measure. Now suppose we have a generalized model as defined in Definition 1.1 and the limit measure is a product. We will show that we cannot *standardize* this to standard regular variation on some cone $\mathfrak{C} \subset \mathbb{E}$ ($\mathfrak{C} = \mathbb{E}_\sqcap$ for our case). Recall that when Definition 1.1 holds, we can always standardize Y so in the following we assume Y^* is standardized and only worry about the standardization of X.

Theorem 3.2. *Suppose $X \in \mathbb{R}, Y^* > 0$ are random variables, such that for functions $\alpha(\cdot) > 0, \beta(\cdot) \in \mathbb{R}$, we have*

$$t\mathbf{P}\Big[\Big(\frac{X-\beta(t)}{\alpha(t)}, \frac{Y^*}{t}\Big) \in \cdot\Big] \xrightarrow{v} G \times \nu_1(\cdot) \qquad \text{in } \mathbb{M}_+\big([-\infty,\infty] \times (0,\infty]\big), \tag{3.6}$$

as $t \to \infty$, where $\nu_1(x,\infty] = x^{-1}$, $x > 0$, and G is some finite, non-degenerate distibution on $\mathbb{R}$. Then there does not exist a standardizing function, $f(\cdot)$: Range of $X \to (0,\infty)$, in the sense of Definition 3.1, such that

$$t\mathbf{P}\Big[\Big(\frac{f(X)}{t}, \frac{Y^*}{t}\Big) \in \cdot\Big] \xrightarrow{v} \mu(\cdot) \qquad \text{in } \mathbb{M}_+\big(\mathbb{E}_\sqcap\big), \tag{3.7}$$

where μ satisfies the conditional non-degeneracy conditions.

Proof. Note that Y^* is already standardized here. Suppose there exists a standardization function $f(\cdot)$ such that (3.7) holds. Without loss of generality assume $f(\cdot)$ to be non-decreasing. This implies that for $\mu-$ continuity points (x,y) we have,

$$t\mathbf{P}\Big[\frac{f(X)}{t} \le x, \frac{Y^*}{t} > y\Big] \to \mu\big((-\infty,x] \times (y,\infty]\big) \qquad (t \to \infty)$$

which is equivalent to

$$t\mathbf{P}\Big(\frac{X-\beta(t)}{\alpha(t)} \le \frac{f^{\leftarrow}(xt)-\beta(t)}{\alpha(t)}, \frac{Y^*}{t} > y\Big] \to \mu\big((-\infty,x] \times (y,\infty]\big), \qquad (t \to \infty). \tag{3.8}$$

Since $\mu((-\infty,x] \times (y,\infty]) < \infty$ and is non-degenerate in x, we have as $t \to \infty$ that

$$\frac{f^{\leftarrow}(xt)-\beta(t)}{\alpha(t)} \to h(x) \tag{3.9}$$

for some non-decreasing function $h(\cdot)$ which has at least two points of increase. Thus (3.8) and (3.9) imply that

$$\mu((-\infty,x] \times (y,\infty]) = G(h(x)) \times y^{-1}.$$

Hence $\mu(\cdot)$ turns out to be a product measure which by Lemma 3.1 is not possible. ☐

The final result of this section shows that one can transform from the conditional extreme value model (like Definition 1.1) to the standard model (like equation (3.3)) and vice-versa if and only if the the limit measure in the generalized model is not a product measure.

Proposition 3.3. *We have two parts to this proposition.*

(1) *Suppose we have the conditional extreme value model of Definition 1.1; i.e., we have a random vector $(X,Y) \in \mathbb{R}^2$, and there exists functions $a(t) > 0, b(t) \in \mathbb{R}, \alpha(t) > 0, \beta \in \mathbb{R}$, such that for $\gamma \in \mathbb{R}$,*

$$t\mathbf{P}\Big[\Big(\frac{X-\beta(t)}{\alpha(t)}, \frac{Y-b(t)}{a(t)}\Big)\cdot\Big] \xrightarrow{v} \mu(\cdot), \qquad \text{in } \mathbb{M}_+\big([-\infty,\infty] \times \overline{\mathbb{E}}^{(\gamma)}\big),$$

along with the conditional non-degeneracy conditions (1.4) *and* (1.5). *Hence equation* (1.8) *holds; i.e.,*

$$\lim_{t\to\infty} \frac{\alpha(tc)}{\alpha(t)} = \psi_1(c), \qquad \lim_{t\to\infty} \frac{\beta(tc)-\beta(t)}{\alpha(t)} = \psi_2(c). \tag{3.10}$$

If $(\psi_1,\psi_2)\neq(1,0)$*, then there exists a standardization function* $\boldsymbol{f}=(f_1,f_2)$ *such that* $(X^*,Y^*)=(f_1(X),f_2(Y))$ *is standard regularly varying on* $\mathbb{E}_\sqcap$*; that is*

$$t\mathbf{P}\Big[\Big(\frac{f_1(X)}{t},\frac{f_2(Y)}{t}\Big)\in\cdot\Big]\xrightarrow{v}\mu^{**}(\cdot)\qquad\qquad \textit{in } \mathbb{M}_+\big(\mathbb{E}_\sqcap\big),$$

where μ^{**} *is a non-null Radon measure satisfying the conditional non-degeneracy conditions.*

(2) *Conversely, suppose we have a bivariate random vector* $(X^*,Y^*)\in\mathbb{R}_+^2$ *satisfying*

$$t\mathbf{P}\Big[\Big(\frac{X^*}{t},\frac{Y^*}{t}\Big)\in\cdot\Big]\xrightarrow{v}\mu^{**}(\cdot)\qquad\qquad \textit{in } \mathbb{M}_+\big(\mathbb{E}_\sqcap\big),$$

where μ^{**} *is a non-null Radon measure, satisfying the conditional non-degeneracy conditions. Consider functions* $\alpha(\cdot)>0$, $\beta(\cdot)\in\mathbb{R}$ *such that equation* (3.10) *holds with* $(\psi_1,\psi_2)\neq(1,0)$*. Then there exist functions* $a(\cdot)>0$, $b(\cdot)\in\mathbb{R}$ *satisfying* (1.2) *and* $\lambda(\cdot)\in\mathbb{R}$, $\gamma\in\mathbb{R}$ *such that*

$$t\mathbf{P}\Big[\Big(\frac{\lambda(X^*)-\beta(t)}{\alpha(t)},\frac{b(Y^*)-b(t)}{a(t)}\Big)\in\cdot\ \Big]\xrightarrow{v}\tilde\mu(\cdot)\ \textit{in } \mathbb{M}_+\big([-\infty,\infty]\times\overline{\mathbb{E}}^{(\gamma)}\big),\tag{3.11}$$

where $\tilde\mu$ *is a non-null Radon measure in* $[-\infty,\infty]\times\overline{\mathbb{E}}^{(\gamma)}$ *satisfying the conditional non-degeneracy conditions and* $b(Y^*)\in D(G_\gamma)$*.*

Remark 3.1. The class of limit measures in Definition 1.1 which are not product measures can thus be considered to be obtained from standard regular variation on $\mathbb{E}_\sqcap$ after appropriate marginal transformations.

Proof. (1) This part has been dealt with in Heffernan and Resnick (2007, Section 2.4).

(2) First we simplify the problem. Note that, for (x,y) a continuity point of $\mu(\cdot)$,

$$t\mathbf{P}\Big[\frac{\lambda(X^*)-\beta(t)}{\alpha(t)}\le x,\frac{b(Y^*)-b(t)}{a(t)}>y\Big]\to\tilde\mu\big([-\infty,x]\times(y,\infty]\big)\qquad(t\to\infty)$$

is equivalent to

$$\begin{aligned}t\mathbf{P}\Big(\frac{\lambda(X^*)-\beta(t)}{\alpha(t)}\le x,\frac{Y^*}{t}>y\Big)&\to\tilde\mu\big([-\infty,x]\times(h(y),\infty]\big)\qquad(t\to\infty)\\&=:\mu^*\big([-\infty,x]\times(y,\infty]\big)\end{aligned}\tag{3.12}$$

where

$$h(y)=\begin{cases}(1+\gamma y)^{\frac{1}{\gamma}} & \gamma\neq0\\ e^y & \gamma=0.\end{cases}\tag{3.13}$$

Hence (3.11) is equivalent to

$$t\mathbf{P}\Big[\Big(\frac{\lambda(X^*)-\beta(t)}{\alpha(t)},\frac{Y^*}{t}\Big)\in\cdot\Big]\xrightarrow{v}\mu^*(\cdot)$$

and μ^* is a non-null Radon measure on $[-\infty,\infty]\times\bar{\mathbb{E}}^{(\gamma)}$ satisfying the conditional non-degeneracy conditions. Hence our proof will show the existence of $\lambda(\cdot)$ satisfying (3.12). Now note that equation (3.10) implies that $\alpha(\cdot)\in RV_\rho$ for some $\rho\in\mathbb{R}$ and $\psi_1(x)=x^\rho$ (see Resnick (2008b, page 14)). The function $\psi_2(\cdot)$ may be identically equal to 0, or

$$\psi_2(x)=\begin{cases}k(x^\rho-1)/\rho, & \text{if } \rho\neq0,x>0\\ k\log x & \text{if } \rho=0,x>0\end{cases}\tag{3.14}$$

for $k\neq0$ (de Haan and Ferreira, 2006, page 373). We have assumed that $(\psi_1,\psi_2)\neq(1,0)$. We will consider three cases: $\rho>0,\rho=0,\rho<0$.

Case 1 : $\rho>0$.

(1) Suppose $\psi_2 \equiv 0$.
Since $\alpha(\cdot) \in RV_\rho$, there exists $\tilde{\alpha}(\cdot) \in RV_\rho$ which is ultimately differentiable and strictly increasing and $\alpha \sim \tilde{\alpha}$ (de Haan and Ferreira, 2006, page 366). Thus $\tilde{\alpha}^{\leftarrow}$ exists. Additionally, we have from Bingham et al. (1987, Theorem 3.1.12(a)), that $\beta(t)/\alpha(t) \to 0$. Hence we have for $x > 0$, as $t \to \infty$,

$$\frac{\tilde{\alpha}(tx) + \beta(t)}{\alpha(t)} = \frac{\tilde{\alpha}(tx)}{\tilde{\alpha}(t)} \cdot \frac{\tilde{\alpha}(t)}{\alpha(t)} + \frac{\beta(t)}{\alpha(t)} \to x^\rho,$$

and inverting we get for $z > 0$

$$\frac{\tilde{\alpha}^{\leftarrow}\big(\alpha(t)z + \beta(t)\big)}{t} \to z^{1/\rho} \qquad (t \to \infty).$$

Thus we have,

$$\begin{aligned} t\mathbf{P}\Big[\frac{\tilde{\alpha}(X^*) - \beta(t)}{\alpha(t)} \le x, \frac{Y^*}{t} > y\Big] &= t\mathbf{P}\Big[\frac{X^*}{t} \le \frac{\tilde{\alpha}^{\leftarrow}(\alpha(t)x + \beta(t))}{t}, \frac{Y^*}{t} > y\Big] \\ &\to \mu^{**}\big([0, x^{1/\rho}] \times (y, \infty]\big). \end{aligned}$$

Set $\lambda(\cdot) = \tilde{\alpha}(\cdot)$ and this defines $\tilde{\mu}$.

(2) Now suppose $\psi_2 \neq 0$.
Therefore

$$\psi_2(x) = \lim_{t\to\infty} \frac{\beta(tx) - \beta(t)}{\alpha(t)} = k(x^\rho - 1)/\rho;$$

that is, $\beta(\cdot) \in RV_\rho$ and $k > 0$. There exists $\tilde{\beta}$ which is ultimately differentiable and strictly increasing and $\tilde{\beta} \sim \beta$ (de Haan and Ferreira, 2006, page 366). Thus $\tilde{\beta}^{\leftarrow}$ exists. Then we have for $x > 0$, as $t \to \infty$,

$$\begin{aligned} \frac{\tilde{\beta}(tx) - \beta(t)}{\alpha(t)} &= \frac{\tilde{\beta}(tx) - \beta(tx)}{\alpha(t)} + \frac{\beta(tx) - \beta(t)}{\alpha(t)} \\ &= \frac{\tilde{\beta}(tx) - \beta(tx)}{\beta(tx)} \frac{\beta(tx)}{\alpha(tx)} \frac{\alpha(tx)}{\alpha(t)} + \frac{\beta(tx) - \beta(t)}{\alpha(t)} \\ &\to (1-1) \cdot \frac{1}{\rho} \cdot x^\rho + k\frac{x^\rho - 1}{\rho} = k\frac{x^\rho - 1}{\rho}. \end{aligned}$$

Inverting, we get as $t \to \infty$,

$$\frac{\tilde{\beta}^{\leftarrow}\big(\alpha(t)x + \beta(t)\big)}{t} \overset{t\to\infty}{\to} (1 + \frac{\rho x}{k})^{1/\rho}.$$

Thus we have,

$$\begin{aligned} t\mathbf{P}\Big[\frac{\tilde{\beta}(X^*) - \beta(t)}{\alpha(t)} \le x, \frac{Y^*}{t} > y\Big] &= t\mathbf{P}\Big[\frac{X^*}{t} \le \frac{\tilde{\beta}^{\leftarrow}(\alpha(t)x + \beta(t))}{t}, \frac{Y^*}{t} > y\Big] \\ &\to \mu^{**}\big([0, (1 + \frac{\rho x}{k})^{1/\rho}] \times (y, \infty]\big). \end{aligned}$$

Here we can set $\lambda(\cdot) = \tilde{\beta}(\cdot)$ and this defines $\tilde{\mu}$.

Case 2: $\rho = 0$.

We have $\psi_1(x) = 1, \psi_2(x) = k \log x$ for $x > 0$ and some $k \in \mathbb{R}$. By assumption, $(\psi_1, \psi_2) \neq (1, 0)$ and hence $k \neq 0$. First assume that $k > 0$, which means $\beta \in \Pi_+(\alpha)$. From property (2) for π-varying functions (Section 6.1), there exists $\tilde{\beta}(\cdot)$ which is continuous, strictly increasing and $\beta - \tilde{\beta} = o(\alpha)$. If $\beta(\infty) = \tilde{\beta}(\infty) = \infty$, we have for $x > 0$,

$$\begin{aligned} \frac{\tilde{\beta}(tx) - \beta(t)}{\alpha(t)} &= \frac{\tilde{\beta}(tx) - \beta(tx)}{\alpha(tx)} \frac{\alpha(tx)}{\alpha(t)} + \frac{\beta(tx) - \beta(t)}{\alpha(t)} \\ &\to 0 + k \log x, \end{aligned}$$

and inverting, we get for $z \in \mathbb{R}$, as $t \to \infty$,

$$\frac{\tilde{\beta}^{\leftarrow}\big(\alpha(t)z + \beta(t)\big)}{t} \to e^{z/k}.$$

Thus we have,

$$\begin{aligned} t\mathbf{P}\Big(\frac{\tilde{\beta}(X^*) - \beta(t)}{\alpha(t)} \le x, \frac{Y^*}{t} > y\Big) &= t\mathbf{P}\Big(\frac{X^*}{t} \le \frac{\tilde{\beta}^{\leftarrow}(\alpha(t)x + \beta(t))}{t}, \frac{Y^*}{t} > y\Big) \\ &\to \mu\big([0, e^{k/x}] \times (y, \infty]\big). \end{aligned}$$

If $\beta(\infty) = \tilde{\beta}(\infty) = B < \infty$, define

$$\beta^*(t) = \frac{1}{B - \tilde{\beta}(t)}, \ \alpha^*(t) = \frac{\alpha(t)}{(B - \tilde{\beta}(t))^2}$$

and from property 3(b), for π-varying functions in Section 6.1, we have that $\beta^* \in \Pi_+(\alpha^*)$, $\beta^*(t) \to \infty$ and $\frac{B-\tilde{\beta}(t)}{\alpha(t)} \overset{t\to\infty}{\to} \infty$. Hence we have reduced to the previous case which implies,

$$t\mathbf{P}\Big(\frac{\beta^*(X^*) - \beta^*(t)}{\alpha^*(t)} \le x, \frac{Y^*}{t} > y\Big) \to \mu\big([0, e^{k/x}] \times (y, \infty]\big).$$

This is equivalent to

$$t\mathbf{P}\Big(\frac{\tilde{\beta}(X^*) - \tilde{\beta}(t)}{\alpha(t)} \le \frac{x}{1 + \frac{\alpha(t)x}{B-\tilde{\beta}(t)}}, \frac{Y^*}{t} > y\Big) \to \mu\big([0, e^{k/x}] \times (y, \infty]\big)$$

and since $\frac{B-\tilde{\beta}(t)}{\alpha(t)} \overset{t\to\infty}{\to} \infty$ implies $\frac{\alpha(t)}{B-\tilde{\beta}(t)} \overset{t\to\infty}{\to} 0$, we can write

$$t\mathbf{P}\Big(\frac{\tilde{\beta}(X^*) - \tilde{\beta}(t)}{\alpha(t)} \le x, \frac{Y^*}{t} > y\Big) \to \mu\big([0, e^{k/x}] \times (y, \infty]\big)$$

which implies that

$$t\mathbf{P}\Big(\frac{\tilde{\beta}(X^*) - \beta(t)}{\alpha(t)} \le x, \frac{Y^*}{t} > y\Big) \to \mu\big([0, e^{k/x}] \times (y, \infty]\big) \qquad \text{since, } \beta - \tilde{\beta} = o(\alpha)$$

and we have produced the required transformation $\lambda(\cdot) = \tilde{\beta}(\cdot)$.

The case for which $k < 0$; i.e., $\beta \in \Pi_-(\alpha)$ can be proved similarly.

Case 3 : $\rho < 0$. This case is similar to the case for $\rho > 0$.

(1) Suppose $\psi_2 \equiv 0$.
Since $\alpha(\cdot) \in RV_\rho$, there exists $\tilde{\alpha}(\cdot) \in RV_\rho$ which is ultimately differentiable and strictly decreasing and $\alpha \sim \tilde{\alpha}$ (de Haan and Ferreira, 2006, page 366). Thus $\tilde{\alpha}^{\leftarrow}$ exists. Additionally, we have from Bingham et al. (1987), Theorem 3.1.10(a),(c) that $\beta(\infty) := \lim_{t\to\infty} \beta(t)$ exists and is finite, and $(\beta(\infty) - \beta(t))/\alpha(t) \to 0$.

$$\frac{\tilde{\alpha}(tx) + \beta(t) - \beta(\infty)}{\alpha(t)} = \frac{\tilde{\alpha}(tx)}{\tilde{\alpha}(t)} \cdot \frac{\tilde{\alpha}(t)}{\alpha(t)} + \frac{\beta(t) - \beta(\infty)}{\alpha(t)} \overset{t\to\infty}{\to} x^\rho,$$

inverting which we get, for $z > 0$, as $t \to \infty$,

$$\frac{\tilde{\alpha}^{\leftarrow}\big(\alpha(t)z + \beta(\infty) - \beta(t)\big)}{t} \to z^{1/\rho}.$$

Thus we have, now taking $x < 0$,

$$t\mathbf{P}\Big(\frac{\beta(\infty) - \tilde{\alpha}(X^*) - \beta(t)}{\alpha(t)} \le x, \frac{Y^*}{t} > y\Big) = t\mathbf{P}\Big(\frac{X^*}{t} \ge \frac{\tilde{\alpha}^{\leftarrow}(-\alpha(t)x + \beta(\infty) - \beta(t))}{t}, \frac{Y^*}{t} > y\Big)$$

$$\to \mu\big([(-x)^{1/\rho}, \infty] \times (y, \infty]\big) =: \tilde{\mu}\big([-\infty, x] \times (y, \infty]\big).$$

Therefore we can set $\lambda(\cdot) = \beta(\infty) - \tilde{\alpha}(\cdot)$.

(2) Now suppose $\psi_2 \neq 0$.
Therefore $\psi_2(x) = \lim\limits_{t\to\infty} \frac{\beta(tx)-\beta(t)}{\alpha(t)} = k(x^\rho - 1)/\rho$; i.e., $\beta(\cdot) \in RV_\rho$ and $k < 0$. There exists $\tilde{\beta} \in RV_\rho$ which is ultimately differentiable and strictly decreasing and $\tilde{\beta} \sim \beta$ (de Haan and Ferreira, 2006, page 366). Thus $\tilde{\beta}^{\leftarrow}$ exists. We also have $\beta(\infty) := \lim_{t\to\infty} \beta(t)$ exists and is finite, and $(\beta(\infty) - \beta(t))/\alpha(t) \to \frac{k}{|\rho|}$ (de Haan and Ferreira, 2006, page 373). Then we have for $x > 0$, as $t \to \infty$,

$$\frac{\tilde{\beta}(tx) - \beta(t)}{\alpha(t)} \to k\frac{x^\rho - 1}{\rho},$$

inverting which we get, as $t \to \infty$,

$$\frac{\tilde{\beta}^{\leftarrow}\big(\alpha(t)x + \beta(t)\big)}{t} \to (1 + \frac{\rho x}{k})^{1/\rho}.$$

Thus we have,

$$t\mathbf{P}\Big(\frac{\tilde{\beta}(X^*) - \beta(t)}{\alpha(t)} \leq x, \frac{Y^*}{t} > y\Big) = t\mathbf{P}\Big(\frac{X^*}{t} \leq \frac{\tilde{\beta}^{\leftarrow}(\alpha(t)x + \beta(t))}{t}, \frac{Y^*}{t} > y\Big)$$
$$\to \mu\big([0, (1 + \frac{\rho x}{k})^{1/\rho}] \times (y, \infty]\big).$$

Here we can set $\lambda(\cdot) = \tilde{\beta}(\cdot)$.

□

Remark 3.2. Suppose that we have

$$t\mathbf{P}\Big[\Big(\frac{X - \beta(t)}{\alpha(t)}, \frac{Y^*}{t}\Big) \in \cdot\Big] \xrightarrow{v} \mu^*(\cdot) = H \times \nu_1(\cdot),$$

in $\mathbb{M}_+([-\infty,\infty] \times (0,\infty])$. We are assuming $\mu^*(\cdot)$ is a product measure. Let

$$X^* = \frac{(X - \beta(Y^*))Y^*}{\alpha(Y^*)}.$$

Then for continuity points (x, y) of the limit

$$t\mathbf{P}\Big(\frac{X^*}{t} \leq x, \frac{Y^*}{t} > y\Big) \to \int_0^{1/y} H(xv)dv$$

in $\mathbb{M}_+([-\infty,\infty] \times (0,\infty])$. It is easy to check that the limit measure is homogeneous of order -1. Thus, a standardization of (X, Y^*) exists even when we have a limit measure which is a product. Note this standardization is not in the sense of Definition 3.1, and it represents a change of co-ordinate system which is more complex than just a marginal transformation.

3.3. A characterization of regular variation on $\mathbb{E}_\sqcap$.

Standard regular variation on $\mathbb{E}$ was characterized by de Haan (1978) in terms of one dimensional regular variation of max linear combinations and Resnick (2002) provides a characterization of hidden regular variation in $\mathbb{E}$ and $\mathbb{E}_0$ in terms of max and min linear combinations of the random vector respectively. We provide a result in the same spirit for regular variation on $\mathbb{E}_\sqcap$.

Proposition 3.4. *Suppose $(X, Y) \in \mathbb{R}^2$ is a random vector and $\mathbf{P}(X = 0) = 0$. Then the following are equivalent:*

1. *(X, Y) is standard multivariate regularly varying on $\mathbb{E}_\sqcap$ with a limit measure satisfying the non-degeneracy conditions (1.4) and (1.5).*
2. *For all $a \in (0, \infty]$ we have*

$$\lim_{t\to\infty} tP\Big(\frac{\min(aX, Y)}{t} > y\Big) = c(a)y^{-1}, \qquad y > 0,$$

for some non-constant, non-decreasing function $c : (0, \infty] \to (0, \infty)$.

Proof. Since $\mathbf{P}(X = 0) = 0$ we have $\mathbf{P}(X > 0) = 1$.

(2) $\Rightarrow$ (1) : Assume that

$$\lim_{t\to\infty} tP\Big(\frac{\min(aX, Y)}{t} > y\Big) = c(a)y^{-1}, \qquad y > 0,$$

for some function $c : (0, \infty] \to (0, \infty)$. Then for $x \geq 0, y > 0$,

$$\begin{aligned} t\mathbf{P}\Big(\frac{X}{t} \leq x, \frac{Y}{t} > y\Big) &= t\mathbf{P}\Big(\frac{Y}{t} > y\Big) - t\mathbf{P}\Big(\frac{X}{t} > x, \frac{Y}{t} > y\Big) \\ &= t\mathbf{P}\Big(X > 0, \frac{Y}{t} > y\Big) - t\mathbf{P}\Big(\frac{(y/x)X}{t} > y, \frac{Y}{t} > y\Big) \\ &= t\mathbf{P}\Big(\frac{\min(a_1X, Y)}{t} > y\Big) - t\mathbf{P}\Big(\frac{\min((y/x)X, Y)}{t} > y\Big) \qquad \text{where } a_1 = \infty \\ &\to c(\infty)y^{-1} - c(y/x)y^{-1} =: \nu([0, x] \times (y, \infty]). \end{aligned}$$

Since $c(\cdot)$ is non-decreasing and non-constant, ν is a non-null Radon measure on $\mathbb{E}_\sqcap$ and we have our result. The non-degeneracy of ν follows from the fact that $c(\cdot)$ is a non-constant function.

(1) $\Rightarrow$ (2) : Now assume that (X, Y) is standard multivariate regularly varying on $\mathbb{E}_\sqcap$. Hence there exists a non-degenerate Radon measure ν on $\mathbb{E}_\sqcap$ such that

$$\lim_{t\to\infty} t\mathbf{P}\Big(\frac{X}{t} \leq x, \frac{Y}{t} > y\Big) = \nu([0, x] \times (y, \infty]),$$

and for any $a \in (0, \infty]$

$$\begin{aligned} t\mathbf{P}\Big(\frac{\min(aX, Y)}{t} > y\Big) &= t\mathbf{P}\Big(\frac{X}{t} > \frac{y}{a}, \frac{Y}{t} > y\Big) \\ &= \nu((\frac{y}{a}, \infty] \times (y, \infty]) \\ &= y^{-1}\nu((\frac{1}{a}, \infty] \times (1, \infty]) =: c(a)y^{-1}, \end{aligned}$$

by defining $c(a) = \nu((\frac{1}{a}, \infty] \times (1, \infty])$ and using the homogeneity property (3.16). Note that the conditional non-degeneracy of ν implies that c is non-constant and non-decreasing. Hence we have the result.

□

Remark 3.3. The condition $\mathbf{P}(X = 0)$ can be removed if we assume Y to be heavy-tailed with exponent $\alpha = 1$, i.e., as $t \to \infty$, $t\mathbf{P}(\frac{Y}{t} > y) \to y^{-1}$.

3.4. **Polar co-ordinates.** Proposition 3.3 shows that when the limit measure is not a product measure, we can transform (X, Y) to (X^*, Y^*) such that

$$\mathbf{P}\Big[\Big(\frac{X^*}{t}, \frac{Y^*}{t}\Big) \in \cdot\Big] \xrightarrow{v} \mu^{**}(\cdot) \qquad \text{in } \mathbb{M}_+(\mathbb{E}_\sqcap) \tag{3.15}$$

Hence from Remark 2.2 we have that μ^{**} is homogeneous of order -1:

$$\mu^{**}(cB) = c^{-1}\mu^{**}(B), \quad \forall c > 0, B \in \mathcal{E}_\sqcap. \tag{3.16}$$

Hence μ^{**} has a spectral form. Further discussion on the spectral form is available in Heffernan and Resnick (2007, Section 3.2). We provide a few facts here. For convenience let us take the norm

$$\|(x,y)\| = |x| + |y|, \quad (x,y) \in \mathbb{R}^2,$$

although any other norm would work too. Now, the standard argument using homogeneity (Resnick, 2008b, Chapter 5) yields for $r > 0$ and Λ a Borel subset of $[0,1)$,

$$\begin{aligned}&\mu^{**}\Big\{(x,y) \in [0,\infty] \times (0,\infty] : x+y > r, \frac{x}{x+y} \in \Lambda\Big\}\\ &= r^{-1}\mu^{**}\Big\{(x,y) \in [0,\infty] \times (0,\infty] : x+y > 1, \frac{x}{x+y} \in \Lambda\Big\} =: r^{-1}S(\Lambda).\end{aligned} \tag{3.17}$$

where S is a Radon measure on $[0,1)$. Note that from (3.17), we can calculate for $x > 0, y > 0$,

$$\mu^{**}\big([0,x] \times (y,\infty]\big) = y^{-1}\int_0^{x/(x+y)} (1-w)S(dw) - x^{-1}\int_0^{x/(x+y)} wS(dw). \tag{3.18}$$

S need not be a finite measure on $[0,1)$ but to guarantee that

$$H^{**}(x) := \mu^{**}\big([0,x] \times (1,\infty]\big) \tag{3.19}$$

is a probability measure, we can see by taking $x \to \infty$ in (3.18) that we need

$$\int_0^1 (1-w)S(dw) = 1. \tag{3.20}$$

Conclusion: The class of limits μ^{**} in (3.15) or conditional limits

$$H^{**}(x) = \lim_{t\to\infty} P\Big[\frac{X^*}{t} \le x | Y > t\Big]$$

is indexed by Radon measures S on $[0,1)$ satisfying the integrability condition (3.20).

Example 3.1 (Finite angular measure)**.** Suppose S is uniform on $[0,1)$, $S(dw) = 2dw$, so that equation (3.20) is satisfied. Then we have

$$\mu^{**}([0,x] \times (y,\infty]) = \frac{x}{y(x+y)}. \tag{3.21}$$

Putting $y = 1$ we get that

$$H^{**}(x) = 1 - \frac{1}{1+x}, \quad x > 0,$$

which is a Pareto distribution.

Example 3.2 (Infinite angular measure)**.** Now suppose S has angular measure $S(dw) = \frac{1}{1-w}dw$. This also satisfies equation (3.20). Now we have

$$\mu^{**}([0,x] \times (y,\infty]) = \frac{1}{y} + \frac{1}{x}\log(1 - \frac{x}{x+y}). \tag{3.22}$$

Putting $y = 1$ we get that

$$H^{**}(x) = 1 - \frac{1}{x}\log(1+x), \quad x > 0.$$

Here H^{**} is continuously increasing, $\lim_{x\downarrow 0} H^{**}(x) = 0$ and $\lim_{x\uparrow\infty} H^{**}(x) = 1$, and hence $H^{**}(\cdot)$ is a valid distribution function. Note that S has infinite angular measure. One way to get infinite angular measures satisfying (3.20) is to take $S(dw) = \frac{1}{1-w}F(dw)$ for probability measures $F(\cdot)$ on $[0,1)$.

4. Extending the Conditional Extreme Value model to a Multivariate Extreme Value Model

Observe that the CEVM assumes the existence of a vague limit in a subset of the Euclidean space which is smaller than that in case of classical MEVT. So it is natural to ask when can we extend a CEVM to a MEVT model. We answer this question in the current section. Clearly, any extension of the CEVM to MEVT will require X to also have a distribution in a domain of attraction. The first Proposition provides a sufficient condition for such an extension.

Proposition 4.1. *Suppose we have $(X,Y) \in \mathbb{R}^2$ and non-negative functions $\alpha(\cdot)$, $a(\cdot)$ and real functions $\beta(\cdot)$, $b(\cdot)$ such that*

$$t\mathbf{P}\Big[\Big(\frac{X-\beta(t)}{\alpha(t)}, \frac{Y-b(t)}{a(t)}\Big) \in \cdot\Big] \xrightarrow{v} \mu(\cdot) \text{ in } \mathbb{M}_+\big([-\infty,\infty] \times \overline{\mathbb{E}}^{(\gamma)}\big),$$

for some $\gamma \in \mathbb{R}$ where μ satisfies the appropriate conditional non-degeneracy conditions corresponding to (1.4)–(1.6). Also assume that $X \in D(G_\lambda)$ for some $\lambda \in \mathbb{R}$; i.e., there exists functions $\chi(t) > 0, \phi(t) \in \mathbb{R}$ such that for continuity points $x \in \mathbb{E}^{(\lambda)}$ of the limit G_λ we have

$$t\mathbf{P}\Big(\frac{X-\phi(t)}{\chi(t)} > x\Big) \to (1+\lambda x)^{-1/\lambda},\ 1+\lambda x > 0.$$

If $\lim_{t\to\infty} \chi(t)/\alpha(t)$ exists and is in $(0,\infty]$ then (X,Y) is in the domain of attraction of a multivariate extreme value distribution on $\mathbb{E}^{(\lambda,\gamma)}$; that is,

$$t\mathbf{P}\Big[\Big(\frac{X-\phi(t)}{\chi(t)}, \frac{Y-b(t)}{a(t)}\Big) \in \cdot\Big] \xrightarrow{v} (\mu \diamond \nu)(\cdot) \qquad \text{in } \mathbb{M}_+\big(\mathbb{E}^{(\lambda,\gamma)}\big)$$

where $(\mu \diamond \nu)(\cdot)$ is a Radon measure on $\mathbb{E}^{(\lambda,\gamma)}$.

Proof. The proof is a consequence of cases 1 and case 2 of Theorem 2.2. □

In the next result we characterize extension of CEVM to MEVT in terms of polar co-ordinates. Assume that (X,Y) can be standardized to (X^*,Y^*) which is regularly varying on $\mathbb{E}_\sqcap$. The following result provides a sufficient condition for an extension of regular variation on $\mathbb{E}_\sqcap$ to an asymptotically tail equivalent regular variation on $\mathbb{E}$. A short discussion on multivariate tail equivalence is provided in the appendix in Section 6.2.

Proposition 4.2. *Suppose $(X,Y) \in \mathbb{R}^2$ is standard regularly varying on the cone $\mathbb{E}_\sqcap$ with limit measure $\nu_\sqcap$ and angular measure $S_\sqcap$ on $[0,1)$. Then the following are equivalent.*

1. *$S_\sqcap$ is finite on $[0,1)$.*
2. *There exists a random vector (X^*,Y^*) defined on $\mathbb{E}$ such that*

$$(X^*,Y^*) \overset{te(\mathbb{E}_\sqcap)}{\sim} (X,Y)$$

and (X^*,Y^*) is multivariate regularly varying on $\mathbb{E}$ with limit measure ν such that $\nu|_{\mathbb{E}_\sqcap} = \nu_\sqcap$.

Proof. Consider each implication separately.

(1) ⇒ (2) : Define the polar coordinate transformation $(R,\Theta) = (X+Y, \frac{X}{X+Y})$. From Section 3.4 we have any $r > 0$, and Λ a Borel subset of $[0,1)$, as $t \to \infty$,

$$t\mathbf{P}\Big[\frac{R}{t} > r, \Theta \in \Lambda\Big] \to r^{-1} S_\sqcap(\Lambda).$$

Note equation (3.17) implies that the right side in the previous line is also equal to

$$\nu_\sqcap\{(x,y) \in \mathbb{E}_\sqcap : x+y > r, \frac{x}{x+y} \in \Lambda\}.$$

Since $S_\sqcap$ is finite on [0,1), the distribution of Θ is finite on $[0,1)$. Assume $S[0,1) = 1$ so that it is a probability measure and extend the measure $S_\sqcap$ to $[0,1]$ by putting $S_\sqcap(\{1\}) = 0$. Let us define R_0

and Θ_0 such that they are independent, Θ_0 has distribution given by the extended $S_{\sqcap}$ on [0,1] and R_0 has the standard Pareto distribution. Define

$$(X^*, Y^*) = (R_0\Theta_0, R_0(1-\Theta_0)).$$

Clearly (X^*, Y^*) is regularly varying on $\mathbb{E}$, now with the standard scaling and limit measure ν where $\nu|_{\mathbb{E}_{\sqcap}} = \nu_{\sqcap}$.

(2) ⇒ (1) : Referring to (3.17) note that

$$S_{\sqcap}\big([0,1)\big) = \nu_{\sqcap}\{(x,y) \in \mathbb{E}_{\sqcap} : x+y > 1\}.$$

Since (X^*, Y^*) is regularly varying on $\mathbb{E}$, we have

$$t\mathbf{P}(\frac{R}{t} > 1) = t\mathbf{P}(\frac{X+Y}{t} > 1) \to \nu\{(x,y) \in \mathbb{E}_{\sqcap} : x+y > 1\} < \infty.$$

But

$$\nu\{(x,y) \in \mathbb{E}_{\sqcap} : x+y>1\} = \nu_{\sqcap}\{(x,y) \in \mathbb{E}_{\sqcap} : x+y>1\} = S_{\sqcap}\big([0,1)\big).$$

Hence $S_{\sqcap}$ is finite on $[0,1)$.

□

5. Examples

In this section we look at examples which help us understand how the conditional model differs from the usual multivariate extreme value model.

Example 5.1. We start by considering the 2-dimensional non-negative orthant. This example emphasizes the fact that we need different normalizations for different cones. This is known for hidden regular variation with the cones $\mathbb{E}$ and $\mathbb{E}_0$ (Example 5.1 in Maulik and Resnick (2005)). We still need a different normalization for the cones $\mathbb{E}_{\sqcap}$ and $\mathbb{E}_{\sqsupset}$.

Let X and Z be i.i.d. *Pareto*(1) random variables. Define $Y = X^2 \wedge Z^2$. Then it is easy to see that the following hold:

(i) In $\mathbb{M}_+(\mathbb{E})$

$$t\mathbf{P}\Big[\Big(\frac{X}{t}, \frac{Y}{t}\Big) \in ([0,x]\times[0,y])^c\Big] \to \frac{1}{x} + \frac{1}{y}, \quad x \vee y > 0. \tag{5.1}$$

(ii) In $\mathbb{M}_+(\mathbb{E}_0)$: For $\frac{1}{2} < \alpha < 1$,

$$t\mathbf{P}\Big[\Big(\frac{X}{t^{\alpha}}, \frac{Y}{t^{2(1-\alpha)}}\Big) \in (x,\infty]\times(y,\infty]\Big] \to \frac{1}{x\sqrt{y}}, \quad x \wedge y > 0,$$

or in standard form,

$$t\mathbf{P}\Big[\Big(\frac{X^{1/\alpha}}{t}, \frac{Y^{1/2(1-\alpha)}}{t}\Big) \in (x,\infty]\times(y,\infty]\Big] \to \frac{1}{x^{\alpha}y^{1-\alpha}}, \quad x \wedge y > 0, \tag{5.2}$$

(iii) In $\mathbb{M}_+(\mathbb{E}_{\sqcap})$, the limit is not a product measure,

$$t\mathbf{P}\Big[\Big(\frac{X}{t^{1/2}}, \frac{Y}{t}\Big) \in [0,x]\times(y,\infty]\Big] \to \frac{1}{y} - \frac{1}{\sqrt{y}} \times \frac{1}{x \vee \sqrt{y}}, \quad x \geq 0, y > 0,$$

so a standard form exists,

$$t\mathbf{P}\Big[\Big(\frac{X^2}{t}, \frac{Y}{t}\Big) \in [0,x]\times(y,\infty]\Big] \to \frac{1}{y} - \frac{1}{\sqrt{y}} \times \frac{1}{\sqrt{x} \vee \sqrt{y}}, \quad x \geq 0, y > 0. \tag{5.3}$$

(iv) In $\mathbb{M}_+(\mathbb{E}_\sqsupset)$, again, the limit is not a product measure,

$$t\mathbf{P}\Big[\Big(\frac{X}{t},\frac{Y}{t^2}\Big)\in(x,\infty]\times[0,y]\Big]\to\frac{1}{x}-\frac{1}{x\vee\sqrt{y}},\quad x>0,y\geq 0,$$

so a standard form exists,

$$t\mathbf{P}\Big[\Big(\frac{X}{t},\frac{Y^{1/2}}{t}\Big)\in(x,\infty]\times[0,y]\Big]\to\frac{1}{x}-\frac{1}{x\vee y},\quad x>0,y\geq 0. \tag{5.4}$$

These results can also be viewed in terms of polar co-ordinates by using the transformation $(r,\theta):(x,y)\mapsto(x+y,\frac{x}{x+y})$ (Section 3.4). Note that the absolute value of the Jacobian of the inverse transformation here is $|J|=r$. Hence,

$$f_{R,\Theta}(r,\theta)=rf_{X,Y}(r\theta,r(1-\theta)).$$

Let us look at the different cones in cases (i)–(iii).

(i) The angular measure has a point mass at 0 and 1,

$$S(d\theta)=\delta_{\{0\}}(d\theta)+\delta_{\{1\}}(d\theta).$$

(ii) The limit measure in standard form is $\mu((x,\infty]\times(y,\infty])=\frac{1}{x^\alpha y^{1-\alpha}},\quad x\wedge y>0.$ for $\frac{1}{2}<\alpha<1$. Hence,

$$\mu'(x,y)=\frac{\partial}{\partial x}\frac{\partial}{\partial y}\mu((x,\infty]\times(y,\infty])=\frac{\alpha(1-\alpha)}{x^{\alpha+1}y^{2-\alpha}}.$$

Taking the polar coordinate transformation

$$\mu'_{R,\Theta}(r,\theta)=r\frac{\alpha(1-\alpha)}{(r\theta)^{\alpha+1}(r(1-\theta))^{2-\alpha}}=r^{-2}\frac{\alpha(1-\alpha)}{\theta^{\alpha+1}(1-\theta)^{2-\alpha}}.$$

The right side is a product, as expected. Thus the angular measure has density

$$S(d\theta)=\frac{\alpha(1-\alpha)}{\theta^{\alpha+1}(1-\theta)^{2-\alpha}}\mathbf{1}_{\{0<\theta<1\}}d\theta.$$

(iii) The limit measure in standard form is

$$\mu([0,x]\times(y,\infty])=\frac{1}{y}-\frac{1}{\sqrt{y}}\times\frac{1}{\sqrt{x}\vee\sqrt{y}},\quad x\geq 0,y>0,$$

which is equivalent to

$$\mu((x,\infty]\times(y,\infty])=\frac{1}{\sqrt{y}}\times\frac{1}{\sqrt{x}\vee\sqrt{y}},\quad x\geq 0,y>0,$$

Hence, for $x>y>0$

$$\mu'(x,y)=\frac{\partial}{\partial x}\frac{\partial}{\partial y}\mu((x,\infty]\times(y,\infty])=\frac{1}{4}\frac{1}{x^{3/2}y^{3/2}}.$$

Taking the polar coordinate transformation, we get for $\theta>1/2$,

$$\mu'_{R,\Theta}(r,\theta)=r\frac{1}{4}\frac{1}{(r\theta)^{3/2}(r(1-\theta))^{3/2}}=\frac{1}{4}r^{-2}\frac{1}{\theta^{3/2}(1-\theta)^{3/2}},$$

the density of a product measure. For $x\leq y$ the density does not exist and we have a point mass at $\theta=\frac{1}{2}$ whose weight can be calculated using (3.20). Thus the angular measure has density,

$$S(d\theta)=(2-\sqrt{3})\delta_{\{1/2\}}(d\theta)+\frac{1}{4}\theta^{-3/2}(1-\theta)^{-3/2}\mathbf{1}_{\{1/2<\theta<1\}}d\theta$$

(iv) The angular measure has a point mass at $\frac{1}{2}$,

$$S(d\theta)=2\delta_{\{1/2\}}(d\theta).$$

Example 5.2. Suppose in Definition 1.1 we have functions $\alpha(t) > 0, \beta(t) = 0$ such that

$$\lim_{t\to\infty} \frac{\alpha(tc)}{\alpha(t)} = \psi_1(c) = c^\rho, \qquad \lim_{t\to\infty} \frac{\beta(tc)-\beta(t)}{\alpha(t)} = \psi_2(c) = 0,$$

for some $\rho \neq 0$. Refer to Heffernan and Resnick (2007, Remark 2, page 545). In such a case, the limit measure μ satisfies:

$$\mu([-\infty, x] \times (y,\infty]) = y^{-1}\mu([-\infty, \frac{x}{y^\rho}] \times (1,\infty]) = y^{-1}H\big(\frac{x}{y^\rho}\big) \tag{5.5}$$

for $x \in \mathbb{R}$ and $y > 0$, where $H(\cdot)$ is a proper non-degenerate distribution. The following is an example of such a limit measure.

Assume $0 < \rho < 1$ and suppose $X \sim Pareto(\rho)$ and $Z \sim Pareto(1-\rho)$ are independent random variables. Define $Y = X \wedge Z$ and we have,

$$\begin{aligned} t\mathbf{P}\Big[\Big(\frac{X}{t}, \frac{Y}{t}\Big) \in [0,x]\times(y,\infty]\Big] &= t\mathbf{P}\Big(\frac{X}{t} \le x, \frac{X}{t} > y, \frac{Z}{t} > y\Big) \\ &= \frac{1}{y^{1-\rho}}\Big(\frac{1}{y^\rho} - \frac{1}{x^\rho}\Big), \qquad \text{(for } x \ge y > 0 \text{ and } t \text{ large)} \\ &= \frac{1}{y}\Big(1 - \frac{y^\rho}{x^\rho}\Big) =: \mu^{**}([-\infty,x]\times(y,\infty]). \end{aligned}$$

Now as in Proposition 3.3, case 1, we have

$$\begin{aligned} t\mathbf{P}\Big[\Big(\frac{\alpha(X)}{\alpha(t)}, \frac{Y}{t}\Big) \in [0,x]\times(y,\infty]\Big] &\to \mu^{**}([0,x^{1/\rho}]\times(y,\infty]) \\ &= \frac{1}{y}\Big(1 - \frac{y^\rho}{x}\Big), \quad x \ge y^\rho > 0 \\ &=: \mu([0,x]\times(y,\infty]). \end{aligned}$$

If we take $H(\cdot)$ to be $Pareto(1)$, then we have the limit measure for $x \ge 0, y > 0$,

$$\mu([-\infty,x]\times(y,\infty]) := \frac{1}{y}H\big(\frac{x}{y^\rho}\big).$$

Example 5.3. This example provides us with a class of limit distributions on $\mathbb{E}_\sqcap$ that can be indexed by distributions on $[0,\infty]$. Suppose R is a Pareto random variable on $[1,\infty)$ with parameter 1 and ξ is a random variable with distribution $G(\cdot)$ on $[0,\infty]$. Assume that ξ and R are independent. Define the bivariate random vector $(X,Y) \in \mathbb{R}_+^2$ as

$$(X,Y) = (R\xi, R).$$

Therefore we have for $y > 0, x \ge 0$ (and $ty > 1$),

$$\begin{aligned} t\mathbf{P}\Big[\frac{X}{t} \le x, \frac{Y}{t} > y\Big] = t\mathbf{P}\Big[\frac{R\xi}{t} \le x, \frac{R}{t} > y\Big] &= t\int_{ty}^{\infty} \mathbf{P}\big[\xi \le \frac{tx}{r}\big]r^{-2}dr \\ &= \int_y^\infty \mathbf{P}\big[\xi \le \frac{x}{s}\big]s^{-2}ds = \int_y^\infty G\big(\frac{x}{s}\big)s^{-2}ds \qquad (\text{ putting } s = \frac{r}{t}) \\ &= \frac{1}{x}\int_0^{x/y} G(s)ds = \mu\big([0,x]\times(y,\infty]\big). \end{aligned}$$

This can be viewed in terms of polar co-ordinates. We know that an angular measure $S(\cdot)$ on $\mathbb{E}_\sqcap$ for $0 \le \eta < 1$ can be given by

$$S([0,\eta]) = \mu\{(u,v) : u+v > 1, \frac{u}{u+v} \le \xi\}.$$

Hence we have

$$\begin{aligned} t\mathbf{P}\Big[\frac{X+Y}{t} > 1, \frac{X}{X+Y} \le \eta\Big] &= t\mathbf{P}\Big[\frac{R\xi + R}{t} > 1, \frac{R\xi}{R\xi + R} \le \eta\Big] \\ &= t\mathbf{P}\Big[\frac{R(1+\xi)}{t} > 1, \xi \le \frac{\eta}{1-\eta}\Big] \\ &= t \int\limits_{0 \le s \le \frac{\eta}{1-\eta}} \mathbf{P}\Big[\frac{R}{t}(1+s) > 1\Big] G(ds) \\ &= t \int\limits_{0 \le s \le \frac{\eta}{1-\eta}} \Big(\frac{t}{1+s} \vee 1\Big)^{-1} G(ds) \\ &= \int\limits_{0 \le s \le \frac{\eta}{1-\eta}} (1+s) G(ds). \qquad \text{for } t > \frac{1}{1-\eta} \end{aligned}$$

But the left side in the previous equation goes to $\mu\{(u,v) : u+v > 1, \frac{y}{u+v} \le \xi\} = S([0,\eta])$ as $t \to \infty$. Hence we have

$$S([0,\eta]) = \int\limits_{0 \le s \le \frac{\eta}{1-\eta}} (1+s) G(ds), \qquad 0 \le \eta < 1.$$

Hence S is a finite angular measure if and only if G has first moment.

6. Appendix

For convenience, this section collects some notation, needed background on regular variation and notions on vague convergence needed for some formulations and proofs.

6.1. Regular variation and the function classes Π.

Regular variation is the mathematical underpinning of heavy tail analysis. It is discussed in many books such as Bingham et al. (1987), de Haan (1970), de Haan and Ferreira (2006), Geluk and de Haan (1987), Resnick (2007, 2008b), Seneta (1976).

A measurable function $U(\cdot) : \mathbb{R}_+ \to \mathbb{R}_+$ is regularly varying at ∞ with index $\rho \in \mathbb{R}$, denoted by $U \in RV_\rho$, if for $x > 0$,

$$\lim_{t\to\infty} \frac{U(tx)}{U(t)} = x^\rho. \tag{6.1}$$

Now let a distribution function $F \in D(G_\gamma)$. For $\gamma > 0$, this entails $\bar{F} = 1 - F \in RV_{-1/\gamma}$ and in general means there exist functions $a(t) > 0, b(t) \in \mathbb{R}$, such that,

$$F^t(a(t)y + b(t)) \to G_\gamma(y), \quad (t \to \infty), \tag{6.2}$$

weakly, where

$$G_\gamma(y) = \exp\{-(1+\gamma y)^{-1/\gamma}\}, \quad 1 + \gamma y > 0,\ \gamma \in \mathbb{R}, \tag{6.3}$$

and the expression on the right is interpreted as $e^{-e^{-y}}$ if $\gamma = 0$. See, for example, Coles (2001), de Haan (1970), Embrechts et al. (1997), Reiss and Thomas (2001), Resnick (2008b). We can and do assume

$$b(t) = \Big(\frac{1}{1 - F(\cdot)}\Big)^{\leftarrow}(t).$$

Thus, we have relation (6.2) is equivalent to

$$t\bar{F}(a(t)y + b(t)) \to (1+\gamma y)^{-1/\gamma}, \quad 1 + \gamma y > 0, \tag{6.4}$$

or taking inverses, as $t \to \infty$,

$$\frac{b(ty) - b(t)}{a(t)} \to \begin{cases} \frac{y^\gamma - 1}{\gamma}, & \text{if } \gamma \neq 0, \\ \log y, & \text{if } \gamma = 0. \end{cases} \tag{6.5}$$

In such a case we say that $b(\cdot)$ is *extended regularly varying* with auxiliary function $a(\cdot)$ and we denote $b \in ERV_\gamma$. In case $\gamma = 0$, we say that $b(\cdot) \in \Pi\big(a(\cdot)\big)$; that is, the function $b(\cdot)$ is Π-*varying* with auxiliary function $a(\cdot)$ (Bingham et al., 1987, de Haan and Ferreira, 2006, Resnick, 2008b).

More generally (de Haan and Ferreira (2006), de Haan and Resnick (1979)) define for an auxiliary function $a(t) > 0$, $\Pi_+(a)$ to be the set of all functions $\pi : \mathbb{R}_+ \mapsto \mathbb{R}_+$ such that

$$\lim_{t\to\infty} \frac{\pi(tx) - \pi(t)}{a(t)} = k \log x, \quad x > 0,\ k > 0. \tag{6.6}$$

The class $\Pi_-(a)$ is defined similarly except that $k < 0$ and

$$\Pi(a) = \Pi_+(a) \cup \Pi_-(a).$$

By adjusting the auxiliary function in the denominator, it is always possible to assume $k = \pm 1$.

Two functions $\pi_i \in \Pi_\pm(a)$, $i = 1, 2$ are $\Pi(a)$-equivalent if for some $c \in \mathbb{R}$

$$\lim_{t\to\infty} \frac{\pi_1(t) - \pi_2(t)}{a(t)} = c.$$

There is usually no loss of generality in assuming $c = 0$. The following are known facts about Π-varying functions.

1. We have $\pi \in \Pi_+(a)$ iff $1/\pi \in \Pi_-(a/\pi^2)$.
2. If $\pi \in \Pi_+(a)$, then (de Haan and Resnick (1979, page 1031) or Bingham et al. (1987, page 159)) there exists a continuous and strictly increasing $\Pi(a)$-equivalent function π_0 with $\pi - \pi_0 = o(a)$.
3. If $\pi \in \Pi_+(a)$, then
$$\lim_{t\to\infty} \pi(t) =: \pi(\infty)$$
exists. If $\pi(\infty) = \infty$, then $\pi \in RV_0$ and $\pi(t)/a(t) \to \infty$. If $\pi(\infty) < \infty$, then $\pi(\infty) - \pi(t) \in \Pi_-(a)$ and $\pi(\infty) - \pi(t) \in RV_0$ and $(\pi(\infty) - \pi(t))/a(t) \to \infty$. (Cf. Geluk and de Haan (1987, page 25).) Furthermore,
$$\frac{1}{\pi(\infty) - \pi(t)} \in \Pi_+\big(a/(\pi(\infty) - \pi(t))^2\big).$$

6.2. **Tail equivalence.** Suppose $\boldsymbol{X}$ and $\boldsymbol{Y}$ are $\mathbb{R}_+^d$-valued random vectors. Then $\boldsymbol{X}$ and $\boldsymbol{Y}$ are *tail equivalent* (in the context of multivariate regular variation) on a cone $\mathfrak{C} \subset \overline{R}_+^d$ if there exists a scaling function $b(t) \uparrow \infty$ such that

$$t\mathbf{P}\Big(\frac{\boldsymbol{X}}{b(t)} \in \cdot\Big) \xrightarrow{v} \nu(\cdot) \qquad \text{and} \qquad t\mathbf{P}\Big(\frac{\boldsymbol{Y}}{b(t)} \in \cdot\Big) \xrightarrow{v} c\nu(\cdot)$$

in $M_+(\mathfrak{C})$ for some $c > 0$ and non-null Radon measure ν on $\mathfrak{C}$. We denote this concept by $X \overset{te(\mathfrak{C})}{\sim} Y$. See Maulik and Resnick (2005).

6.3. **Vague convergence.** For a nice space $\mathbb{E}^*$, that is, a space which is locally compact with countable base (for example, a finite dimensional Euclidean space), denote $\mathbb{M}_+(\mathbb{E}^*)$ for the non-negative Radon measures on Borel subsets of $\mathbb{E}^*$. This space is metrized by the vague metric. The notion of vague convergence in this space is as follows: If $\mu_n \in \mathbb{M}_+(\mathbb{E}^*)$ for $n \geq 0$, then μ_n converge vaguely to μ_0 (written $\mu_n \xrightarrow{v} \mu_0$) if for all bounded continuous functions f with compact support we have

$$\int_{\mathbb{E}^*} f d\mu_n \to \int_{\mathbb{E}^*} f d\mu_0 \quad (n \to \infty).$$

This concept allows us to write (1.2) as

$$tP[\frac{Y - b(t)}{a(t)} \in \cdot] \xrightarrow{v} m_\gamma(\cdot), \tag{6.7}$$

vaguely in $\mathbb{M}_+((-\infty, \infty])$ where

$$m_\gamma((x, \infty]) = (1 + \gamma x)^{-1/\gamma}.$$

Standard references include Kallenberg (1983), Neveu (1977) and Resnick (2008b, Chapter 3).

6.4. **Statements of some Theorems and Proofs:**

6.4.1. *Proof of Theorem 2.1.*

Proof. First, note that if A is relatively compact in $\mathbb{E}_0 = \mathbb{E}_\sqcap \cap \mathbb{E}_\sqsupset$ with $\mu(\partial A) = 0 = \nu(\partial A)$, we will have

$$\mu(A) = \lim_{t\to\infty} t\mathbf{P}\Big[\Big(\frac{X}{t}, \frac{Y}{t}\Big) \in A\Big] = \nu(A) \tag{6.8}$$

(Portmanteau Theorem for vague convergence, Theorem (3.2) in Resnick (2007)). Hence we have a unique vague limit on $M_+(\mathbb{E}_0)$; i.e.,

$$\mu(\cdot)|_{\mathbb{E}_\sqcap \cap \mathbb{E}_\sqsupset} = \nu(\cdot)|_{\mathbb{E}_\sqcap \cap \mathbb{E}_\sqsupset}. \tag{6.9}$$

For $\epsilon > 0$ let us define:

$$B_1^\epsilon = [0, \epsilon) \times [\epsilon, \infty] \in \mathcal{E}_\sqcap,$$
$$B_2^\epsilon = [\epsilon, \infty] \times [0, \infty] \in \mathcal{E}_\sqsupset.$$

We will prove the theorem with the following two claims.

Claim 6.0.1. Let $A \in \mathcal{E}$ be a relatively compact set in $\mathbb{E}$. Define the set

$$[(\mu \diamond \nu)(A)] = \{\mu(A \cap B_1^\epsilon) + \nu(A \cap B_2^\epsilon) : 0 < \epsilon < d(0, A)/\sqrt{2}\}.$$

Then $[(\mu \diamond \nu)(A)]$ is a singleton set and if we denote the unique element of this set by $(\mu \diamond \nu)(A)$, then $(\mu \diamond \nu)(\cdot)$ is a Radon measure on $\mathbb{E}$.

Proof. Since A is relatively compact, $d(0, A) > 0$. Take $0 < \epsilon < \delta < d(0, A)/\sqrt{2}$. Now to show that $[(\mu \diamond \nu)(A)]$ is a singleton, it suffices to check the following:

$$\mu(A \cap B_1^\epsilon) + \nu(A \cap B_2^\epsilon) = \mu(A \cap B_1^\delta) + \nu(A \cap B_2^\delta).$$

Let us define

$$F_1 = B_1^\epsilon \cap B_1^\delta, \quad F_2 = B_1^\epsilon \cap (B_1^\delta)^c, \quad F_3 = (B_1^\epsilon)^c \cap B_1^\delta, \quad F_4 = B_2^\epsilon \cap (B_1^\delta)^c \cap (B_2^\delta)^c, \quad F_5 = B_2^\delta.$$



FIGURE 2. Partitioning $\mathbb{E}_0$

Referring to Figure 2 we have

$$
\begin{aligned}
B_1^\epsilon &= F_1 \cup F_2, & B_1^\delta &= F_1 \cup F_3,\\
B_2^\epsilon &= F_3 \cup F_4 \cup F_5 & B_2^\delta &= F_5.
\end{aligned}
$$

First note that $F_1, F_2, F_3 \in \mathcal{E}_\sqcap$, $F_3, F_4, F_5 \in \mathcal{E}_\sqsupset$. Hence $A \cap F_3 \in \mathcal{E}_0$ and from (6.9) we have

$$\mu(A \cap F_3) = \nu(A \cap F_3)$$

Also note that $\delta < d(0, A)/\sqrt{2}$ implies $A \subset B_1^\delta \cup B_2^\delta = F_1 \cup F_3 \cup F_5$. So

$$A \cap F_2 = \emptyset, A \cap F_4 \ = \emptyset. \tag{6.10}$$

Now

$$
\begin{aligned}
\mu(A \cap B_1^\epsilon) + \nu(A \cap B_2^\epsilon) &= \mu(A \cap (F_1 \cup F_2)) + \nu(A \cap (F_3 \cup F_4 \cup F_5))\\
&= \mu(A \cap F_1) + \mu(A \cap F_2) + \nu(A \cap F_3) + \nu(A \cap F_4) + \nu(A \cap F_5)\\
&= \mu(A \cap F_1) + \mu(A \cap F_3) + \nu(A \cap F_5) \qquad \text{(using (6.9) and (6.10))}\\
&= \mu(A \cap (F_1 \cup F_3)) + \nu(A \cap F_5)\\
&= \mu(A \cap B_1^\delta) + \nu(A \cap B_2^\delta).
\end{aligned}
$$

Thus $[(\mu \diamond \nu)(A)]$ is a singleton and we denote its unique element by $(\mu \diamond \nu)(A)$. It is easy to see that $(\mu \diamond \nu)(\cdot)$ is a Radon measure. □

Claim 6.0.2. Let A be a relatively compact set in $\mathbb{E}$ with $(\mu \diamond \nu)(\partial A) = 0$. Then

$$t\mathbf{P}\Big[\Big(\frac{X}{t}, \frac{Y}{t}\Big) \in A\Big] \to (\mu \diamond \nu)(A).$$

Proof. Choose $0 < \epsilon < d(0, A)$ s.t. $\mu(\partial B_1^\epsilon) = \nu(\partial B_2^\epsilon) = 0$. Now, $\partial(A \cap B_1^\epsilon) \subseteq \partial A \cup \partial B_1^\epsilon$. Therefore

$$
\begin{aligned}
(\mu \diamond \nu)(\partial(A \cap B_1^\epsilon)) &\le (\mu \diamond \nu)(\partial A) + (\mu \diamond \nu)(\partial B_1^\epsilon)\\
&= 0 + \mu(\partial B_1^\epsilon) = 0 \qquad \text{(using Claim 6.0.1 since } \partial B_1^\epsilon \in \mathcal{E}_\sqcap).
\end{aligned}
$$

Thus $A \cap B_1^\epsilon$ is relatively compact in $\mathbb{E}_\sqcap$ and $\mu(\partial(A \cap B_1^\epsilon)) = 0$. Similarly $A \cap B_2^\epsilon$ is relatively compact in $\mathbb{E}_\sqsupset$ and $\nu(\partial(A \cap B_2^\epsilon)) = 0$. Hence we have

$$
\begin{aligned}
t\mathbf{P}\Big[\Big(\frac{X}{t}, \frac{Y}{t}\Big) \in A\Big] &= t\mathbf{P}\Big[\Big(\frac{X}{t}, \frac{Y}{t}\Big) \in A \cap B_1^\epsilon\Big] + t\mathbf{P}\Big[\Big(\frac{X}{t}, \frac{Y}{t}\Big) \in A \cap B_2^\epsilon\Big]\\
&\to \mu(A \cap B_1^\epsilon) + \nu(A \cap B_2^\epsilon)\\
&= (\mu \diamond \nu)(A).
\end{aligned}
$$

□

This claim holds for any $A \in \mathcal{E}$ which is relatively compact with $\mu(\partial A) = 0$. Thus from Portmanteau Theorem for vague convergence (Resnick, 2007, Theorem (3.2)) we have proven Claim 6.0.2.

Now we show that $(\mu \diamond \nu)|_{\mathbb{E}_\sqcap}(\cdot) = \mu(\cdot)$ on $\mathbb{E}_\sqcap$. If $A \in \mathcal{E}_\sqcap$, choose any $0 < \epsilon < d(0, A)/\sqrt{2}$ for defining $(\mu \diamond \nu)(A)$. Now we have $A \cap B_2^\epsilon = A \cap B_2^\epsilon \cap \mathbb{E}_\sqcap \in \mathcal{E}_0$. Therefore

$$
\begin{aligned}
(\mu \diamond \nu)(A) &= \mu(A \cap B_1^\epsilon) + \nu(A \cap B_2^\epsilon)\\
&= \mu(A \cap B_1^\epsilon) + \nu(A \cap B_2^\epsilon \cap \mathbb{E}_\sqcap)
\end{aligned}
$$

and from (6.8), since $A \cap B_2^\epsilon \cap \mathbb{E}_\sqcap \in \mathcal{E}_0$, this is

$$
\begin{aligned}
&= \mu(A \cap B_1^\epsilon) + \mu(A \cap B_2^\epsilon \cap \mathbb{E}_\sqcap)\\
&= \mu(A \cap (B_1^\epsilon \cup (B_2^\epsilon \cap \mathbb{E}_\sqcap))) = \mu(A).
\end{aligned}
$$

We can prove $(\mu \diamond \nu)|_{\mathbb{E}_\sqsupset}(\cdot) = \nu(\cdot)$ on $\mathbb{E}_\sqsupset$ similarly.

□

6.4.2. *Proof of Proposition 2.3.* We will state and prove the result for a simple standard case which is a particularization of Theorem 2.1.

Proposition 6.1. *$(X,Y)\in\mathbb{R}_+^2$ is a bivariate random vector with the following properties:*

1. *(X,Y) has a density $f_{X,Y}(x,y)$.*
2. *Corresponding to regular variation in standardized form we assume,*
$$t^2 f_X(xt)\to x^{-2}, x>1, \qquad t^2 f_Y(yt)\to y^{-2}, y>1. \tag{6.11}$$
3. *The joint density satisfies*
$$t^3 f_{X,Y}(tx,ty)\to g_1(x,y)\in L_1(\mathbb{E}_\sqcap), \tag{6.12}$$
$$t^3 f_{X,Y}(tx,ty)\to g_2(x,y)\in L_1(\mathbb{E}_\sqsupset), \tag{6.13}$$
where $g_1(x,y), g_2(x,y)\geq 0$ are non-trivial, assumed 0 outside of $\mathbb{E}_\sqcap$ and $\mathbb{E}_\sqsupset$ respectively and we have
   - (a) *$v^2 g_1(u,v)$ is a probability density in v for each $u>0$,*
   - (b) *$u^2 g_2(u,v)$ is a probability density in u for each $v>0$.*

Then we have
$$t\mathbf{P}\Big[\Big(\frac{X}{t},\frac{Y}{t}\Big)\in\cdot\Big]\xrightarrow{v}\mu(\cdot) \text{ in } \mathbb{M}_+(\mathbb{E}),$$
where
$$\mu(A)=\int\limits_{\{(u,v)\in A\}} g^*(u,v)dudv \qquad \text{for } A\in\mathcal{E},$$
where $g^*(x,y)=g_1(x,y)\vee g_2(x,y)$.

Remark 6.1. Referring to Theorem 2.1, the conditions in (6.11) correspond to marginal convergences implicit in the Theorem 2.1, and conditions (6.12) and (6.13) correspond to conditions (2.3) and (2.4). Conditions (a) and (b) guarantee that we have proper conditional probability densities.

Proof. Clearly $g^*(\cdot)\in L_1(\mathbb{E})$. Also note that (6.12) and (6.13) imply that
$$g_1(x,y)=g_2(x,y)=g^*(x,y), \qquad \text{for } (x,y)\in\mathbb{E}_0.$$
Hence, for $(u,v)\in\mathbb{E}_0$,
$$\begin{aligned} tf_{\frac{X}{t},\frac{Y}{t}}(u,v) &= t^3 f_{X,Y}(tu,tv)\\ &\to g^*(u,v), \qquad \text{as } t\to\infty, \end{aligned}$$
using (6.12) or (6.13). Now for $x>0, y>0$:
$$\begin{aligned} t\mathbf{P}\Big[\Big(\frac{X}{t},\frac{Y}{t}\Big)\in\big([0,x]\times[0,y]\big)^c\Big] &= t\mathbf{P}\Big(\frac{X}{t}>x\Big)+t\mathbf{P}\Big(\frac{Y}{t}>y\Big)-t\mathbf{P}\Big(\frac{X}{t}>x,\frac{Y}{t}>y\Big)\\ &= t\int\limits_x^\infty f_{\frac{X}{t}}(u)du + t\int\limits_y^\infty f_{\frac{Y}{t}}(v)dv - t\int\limits_x^\infty\int\limits_y^\infty f_{\frac{X}{t},\frac{Y}{t}}(u,v)dudv\\ &\to x^{-1}+y^{-1}-\int\limits_x^\infty\int\limits_y^\infty g^*(u,v)dudv, \qquad \text{as } t\to\infty, \end{aligned}$$
which is a consequence of Scheffe' Theorem. Also note that
$$x^{-1}=\int\limits_x^\infty u^{-2}du=\int\limits_x^\infty u^{-2}\Big(\int\limits_0^\infty u^2 g_2(u,v)dv\Big)du=\int\limits_x^\infty\int\limits_0^\infty g^*(u,v)dvdu,$$

and similarly

$$y^{-1} = \int_0^\infty \int_y^\infty g^*(u,v)dvdu.$$

Therefore as $t \to \infty$,

$$t\mathbf{P}\Big[\Big(\frac{X}{t}, \frac{Y}{t}\Big) \in \big([0,x] \times [0,y]\big)^c\Big] \to \int_x^\infty \int_0^\infty g^*(u,v)dvdu + \int_0^\infty \int_y^\infty g^*(u,v)dvdu - \int_x^\infty \int_y^\infty g^*(u,v)dudv$$

$$= \int_{(u,v)\in([0,x]\times[0,y])^c} g^*(u,v)(u,v)dudv = \mu\big(\big([0,x] \times [0,y]\big)^c\big). \tag{6.14}$$

According to Lemma 6.1, Resnick (2007), proving 6.14 suffices for our proof. □

References

N.H. Bingham, C.M. Goldie, and J.L. Teugels. *Regular Variation.* Cambridge University Press, 1987.

S. G. Coles and J. A. Tawn. Modelling extreme multivariate events. *J. R. Statist. Soc. B*, 53:377–392, 1991.

S.G. Coles. *An Introduction to Statistical Modeling of Extreme Values.* Springer Series in Statistics. London: Springer. xiv, 210 p. , 2001.

Youri Davydov, Ilya Molchanov, and Sergei Zuyev. Stable distributions and harmonic analysis on convex cones. *C. R. Math. Acad. Sci. Paris*, 344(5):321–326, 2007. ISSN 1631-073X.

L. de Haan. *On Regular Variation and Its Application to the Weak Convergence of Sample Extremes.* Mathematisch Centrum Amsterdam, 1970.

L. de Haan and J. de Ronde. Sea and wind: multivariate extremes at work. *Extremes*, 1(1):7–46, 1998.

L. de Haan and A. Ferreira. *Extreme Value Theory: An Introduction.* Springer-Verlag, New York, 2006.

L. de Haan and S.I. Resnick. Limit theory for multivariate sample extremes. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete*, 40:317–337, 1977.

L. de Haan and S.I. Resnick. Conjugate π-variation and process inversion. *Ann. Probab.*, 7(6):1028–1035, 1979. ISSN 0091-1798.

Laurens de Haan. A characterization of multidimensional extreme-value distributions. *Sankhyā Ser. A*, 40(1):85–88, 1978. ISSN 0581-572X.

P. Embrechts, C. Kluppelberg, and T. Mikosch. *Modelling Extreme Events for Insurance and Finance.* Springer-Verlag, Berlin, 1997.

J. L. Geluk and L. de Haan. *Regular Variation, Extensions and Tauberian Theorems*, volume 40 of *CWI Tract*. Stichting Mathematisch Centrum, Centrum voor Wiskunde en Informatica, Amsterdam, 1987. ISBN 90-6196-324-9.

J.E. Heffernan and S.I. Resnick. Hidden regular variation and the rank transform. *Adv. Appl. Prob.*, 37(2):393–414, 2005.

J.E. Heffernan and S.I. Resnick. Limit laws for random vectors with an extreme component. *Ann. Appl. Probab.*, 17 (2):537–571, 2007. ISSN 1050-5164. doi: 10.1214/105051606000000835.

J.E. Heffernan and J.A. Tawn. A conditional approach for multivariate extreme values (with discussion). *JRSS B*, 66(3):497–546, 2004.

O. Kallenberg. *Random Measures.* Akademie-Verlag, Berlin, third edition, 1983. ISBN 0-12-394960-2.

C. Klüppelberg and S. I. Resnick. The Pareto copula, aggregation of risks and the emperor's socks. *Journal of Applied Probability*, 45(1):67–84, 2008.

A.W. Ledford and J.A. Tawn. Statistics for near independence in multivariate extreme values. *Biometrika*, 83(1): 169–187, 1996. ISSN 0006-3444.

A.W. Ledford and J.A. Tawn. Modelling dependence within joint tail regions. *J. Roy. Statist. Soc. Ser. B*, 59(2): 475–499, 1997. ISSN 0035-9246.

A.W. Ledford and J.A. Tawn. Concomitant tail behaviour for extremes. *Adv. in Appl. Probab.*, 30(1):197–215, 1998. ISSN 0001-8678.

K. Maulik and S.I. Resnick. Characterizations and examples of hidden regular variation. *Extremes*, 7(1):31–67, 2005.

T. Mikosch. How to model multivariate extremes if one must? *Statist. Neerlandica*, 59(3):324–338, 2005. ISSN 0039-0402.

T. Mikosch. Copulas: Tales and facts. *Extremes*, 9(1):3–20, 2006. ISSN 1386-1999 (Print) 1572-915X (Online).

J. Neveu. Processus ponctuels. In *École d'Été de Probabilités de Saint-Flour, VI—1976*, pages 249–445. Lecture Notes in Math., Vol. 598, Berlin, 1977. Springer-Verlag.

J. Pickands. Multivariate extreme value distributions. In *43rd Sess. Int. Statist. Inst.*, pages 859–878, 1981.

R.-D. Reiss and M. Thomas. *Statistical Analysis of Extreme Values*. Birkhäuser Verlag, Basel, second edition, 2001. ISBN 3-7643-6487-4.

S.I. Resnick. Hidden regular variation, second order regular variation and asymptotic independence. *Extremes*, 5(4): 303–336, 2002.

S.I. Resnick. *Heavy Tail Phenomena: Probabilistic and Statistical Modeling*. Springer Series in Operations Research and Financial Engineering. Springer-Verlag, New York, 2007. ISBN: 0-387-24272-4.

Sidney I. Resnick. Multivariate regular variation on cones: application to extreme values, hidden regular variation and conditioned limit laws. *Stochastics: An International Journal of Probability and Stochastic Processes*, 80: 269–298, 2008a. ISSN 1744-2508. URL `http://www.informaworld.com/10.1080/17442500701830423`.

Sidney I. Resnick. *Extreme values, regular variation and point processes*. Springer Series in Operations Research and Financial Engineering. Springer, New York, 2008b. ISBN 978-0-387-75952-4. Reprint of the 1987 original.

E. Seneta. *Regularly Varying Functions*. Springer-Verlag, New York, 1976. Lecture Notes in Mathematics, 508.

Bikramjit Das, School of Operations Research and Information Engineering, Cornell University, Ithaca, NY 14853

E-mail address: `bd72@cornell.edu`

Sidney Resnick, School of Operations Research and Information Engineering, Cornell University, Ithaca, NY 14853

E-mail address: `sir1@cornell.edu`